

A Chemical Biology Solution to Problems with Studying Biologically Important but Unstable 9-O-Acetyl Sialic Acids

Zahra Khedri,^{†,||} An Xiao,^{‡,||} Hai Yu,^{‡,||} Corinna Susanne Landig,[†] Wanqing Li,[‡] Sandra Diaz,[†] Brian R. Wasik,[§] Colin R. Parrish,[§] Lee-Ping Wang,^{‡,||} Ajit Varki,^{*,†} and Xi Chen^{*,‡,||}


[†]Glycobiology Research and Training Center, University of California, San Diego, California 92093, United States

[‡]Department of Chemistry, University of California, Davis, California 95616, United States

[§]Department of Microbiology and Immunology, Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, New York 14853, United States

S Supporting Information

ABSTRACT: 9-O-Acetylation is a common natural modification on sialic acids (Sias) that terminate many vertebrate glycan chains. This ester group has striking effects on many biological phenomena, including microbe-host interactions, complement action, regulation of immune responses, sialidase action, cellular apoptosis, and tumor immunology. Despite such findings, 9-O-acetyl sialoglycoconjugates have remained largely understudied, primarily because of marked lability of the 9-O-acetyl group to even small pH variations and/or the action of mammalian or microbial esterases. Our current studies involving 9-O-acetylated sialoglycans on glycan micro-



arrays revealed that even the most careful precautions cannot ensure complete stability of the 9-O-acetyl group. We now demonstrate a simple chemical biology solution to many of these problems by substituting the oxygen atom in the ester with a nitrogen atom, resulting in sialic acids with a chemically and biologically stable 9-N-acetyl group. We present an efficient one-pot multienzyme method to synthesize a sialoglycan containing 9-acetamido-9-deoxy-N-acetylneuraminic acid (Neu5Ac9NAc) and compare it to the one with naturally occurring 9-O-acetyl-N-acetylneuraminic acid (Neu5,9Ac₂). Conformational resemblance of the two molecules was confirmed by computational molecular dynamics simulations. Microarray studies showed that the Neu5Ac9NAc-sialoglycan is a ligand for viruses naturally recognizing Neu5,9Ac₂, with a similar affinity but with much improved stability in handling and study. Feeding of Neu5Ac9NAc or Neu5,9Ac₂ to mammalian cells resulted in comparable incorporation and surface expression as well as binding to 9-O-acetyl-Sia-specific viruses. However, cells fed with Neu5Ac9NAc remained resistant to viral esterases and showed a slower turnover. This simple approach opens numerous research opportunities that have heretofore proved intractable.

The first sialic acid discovered (crystallized by Gunner Blix from a hot mild acid extract of bovine submaxillary mucin in 1936) contained two acetyl groups, only one possibly attached to nitrogen.¹ In retrospect, Blix likely isolated a 9-O-acetyl variant of the common sialic acid N-acetylneuraminic acid (Neu5Ac), 9-O-acetyl-N-acetylneuraminic acid (Neu5,9Ac₂; **1**, Figure 1). Neu5,9Ac₂ along with Neu5Ac and N-glycolylneuraminic acid (Neu5Gc) are the three most frequently occurring sialic acid forms in mammals.² It has




Figure 1. Structures of 9-O-acetyl-N-acetylneuraminic acid (Neu5,9Ac₂; **1**) and its N-acetyl analog Neu5Ac9NAc (**2**).

long been known that 9-O-acetylation of sialic acids masks the recognition by influenza A virus hemagglutinin³ and by other lectins such as Factor H,^{4,5} CD22/Siglec-2,⁵⁻⁷ and Sialoadhesin/Siglec-1,⁷ while being required for the binding of influenza C virus hemagglutinin.^{3,8-11} Despite such early recognition and understanding of their importance, studies of the biological significance of sialoglycans presenting this common O-acetylated form of sialic acid have lagged far behind those of the parent molecules. The reasons are many and have been detailed elsewhere, including their lability to both acidic and basic conditions (often used in standard purification methods for glycans and glycoconjugates), their propensity to migrate from one position to another, and their relative or absolute effects in blocking sialidase action.^{2,3,12} However, when it has

Received: October 21, 2016

Accepted: November 28, 2016

Published: November 28, 2016

been further studied, such O-acetyl group modification turns out to be a key determinant modulating recognition by viruses,^{3,13,14} antibodies,^{15–18} and mammalian lectins,^{6,7,17,19} as well as modulating sialidase action^{2,3,12} and cellular apoptosis.^{20–24}

Overall, exploration of the functions of Neu5,9Ac₂ and its biological and pathological interactions has been greatly hampered by the chemical instability of the O-acetyl group and/or the esterase cleavage of such a group.⁸ Taken together with the fact that it is eliminated by basic conditions during conventional approaches to glycomic analysis such as beta-elimination and permethylation, it has come to the point where 9-O-acetylation tends to be simply ignored in many studies. Synthetic analogs of Neu5,9Ac₂ with chemical modifications of the 9-position of Neu5Ac generated included a 9-N-acetyl analog, 9-acetamido-9-deoxy-N-acetylneuraminic acid (Neu5Ac9NAc; **2**, Figure 1), which was shown to mimic Neu5,9Ac₂ in binding to influenza C virus without being destroyed by esterase activity of the hemagglutinin-esterase.²⁵ This observation was not however further explored for chemical and biological studies of Neu5,9Ac₂. Herein, we report a more efficient chemoenzymatic method for synthesizing Neu5Ac9NAc and a sialoside containing this Neu5,9Ac₂ analog. Applications in glycan microarray and cell feeding studies are presented, laying the foundation for a new approach to elucidate the important roles of 9-O-acetylation of sialic acids.

RESULTS

Problems with Instability of O-Acetyl Sialic Acids in Preparing and Using Glycan Microarrays. The early described glycan microarrays did not have much representation of O-acetyl sialic acids.²⁶ In an attempt to address this deficiency, our groups have been collaborating to synthesize, print, and study matched pairs of sialoglycans that either have or do not have O-acetyl esters. Due to their relative ease of synthesis and handling compared to 7-O- or 8-O-acetyl analogs, sialosides with 9-O-acetylated sialic acids were the ones studied first. Over the past few years, we have been collaborating with several other groups to study the binding of various proteins to such glycans using glycan microarray technology.^{16–19,27–32} In revisiting some of these studies, we noticed that some of the 9-OAc-glycans used for microarray printing were partially de-O-acetylated.

Investigating and Minimizing 9-O-Acetyl Group Loss at Various Steps in Sample Preparation, Preparing and Using of Glycan Microarrays. On the basis of the above observations, we investigated the extent of de-O-acetylation at the final steps of the synthesis and entire glycan arraying process for the model 9-OAc-glycan Neu5,9Ac₂α3Galβ4GlcβProN₃ and its reduced form Neu5,9Ac₂α3Galβ4GlcβProNH₂. The azide-containing glycan was analyzed using nuclear magnetic resonance (NMR) spectrometry and high resolution mass spectrometry (HRMS). It was found to be pure, without any loss of OAc group. The compound was then subjected to hydrogenolysis using H₂ and Pd–C in water with a drop of glacial acetic acid. The reaction was completed in 1 h as detected using MS. To minimize de-O-acetylation during the Celite filtration step which uses Na₂CO₃ as a mild base,³³ filtration after catalytic hydrogenation was replaced by passing the reaction solution through a nylon syringe filter (0.2 μm). The product was then purified using a short C-18 cartridge (water as an eluant) instead of the commonly and previously used Bio-Gel

purification, as it is faster. The fractions containing the pure product were combined and lyophilized. No noticeable loss of the O-acetyl group was observed using the improved procedures described above.

Then, the reduced glycan was quantified for the OAc loss at each step of glycan microarray process. Our results revealed that ~45% of the OAc group was either lost from the 9-position of Neu5Ac or migrated to 7- and 8-positions of the sialic acid upon conventional analysis procedures including release of sialic acids with acetic acid, derivatization with 1,2-diamino-4,5-methylenedioxybenzene (DMB), followed by high performance liquid chromatography (HPLC) analysis of fluorescent adducts (Figure 2A). In comparison, *Arthrobacter ureafaciens* sialidase




Figure 2. DMB derivatization and HPLC analysis of sialic acids released from Neu5,9Ac₂α3Galβ4GlcβProNH₂ using acetic acid (A) and AUS (B).

(AUS)³⁴ treatment followed by DMB derivatization at low temperatures³⁵ showed only 1.5% loss of the OAc group (Figure 2B). This means there is no significant de-O-acetylation during initial preparation of stock solution in water (it should be noted that the final synthetic product is a neutral sodium salt—earlier studies in which the acid form was prepared was associated with some de-O-acetylation, data not shown) and AUS treatment with low temperature DMB derivatization. However, about a 3% loss of OAc was observed during storage in phosphate buffer (pH 8.4) at RT, a condition popular for current microarray printing. Furthermore, about 5% of the sialic acid was de-O-acetylated under the standard blocking condition used for glycan arrays (ethanolamine in Tris-HCl buffer, pH 9.0). The glycan stock solution was also analyzed again after three months of storage at –20 °C and a few freeze–thaw cycles. It was noticed that another 4% of the OAc group was released over three months. Therefore, a small amount of de-O-acetylation occurred under storage conditions and under conditions used for microarray printing and slide blocking. Any further loss of O-acetylation after printing during the binding and analysis, however, could not be directly monitored.

Overall, definitive conclusions about binding of probes can therefore only be made where binding is exclusive to the O-acetylated sialoglycan spots and not to the corresponding non-

O-acetylated ones, as discussed later. A more general conclusion is that, even under the most carefully managed handling by skilled experimentalists, some loss of 9-O-acetylation appears inevitable. Taken together with the knowledge that esterases of bacterial^{36–40} and vertebrate^{41–43} origin are very common in biological systems, there is an intractable problem facing the systematic study of the chemistry and biology of sialic acid 9-O-acetylation.

Chemical Synthesis of Neu5Ac9NAc Precursor 6-Acetamido-6-deoxy-N-acetylmannosamine (ManNAc6-NAc) and Enzymatic Synthesis of Neu5Ac9NAc. A new efficient chemoenzymatic synthetic strategy was developed for the synthesis of Neu5Ac9NAc. The design was to chemically synthesize ManNAc6NAc, a precursor of Neu5Ac9NAc, followed by a sialic acid aldolase-catalyzed reaction^{44,45} for the synthesis of desired Neu5Ac9NAc. ManNAc6NAc was synthesized in 82% yield from previously obtained 6-azido-6-deoxy-N-acetylmannosamine (ManNAc6N₃)⁴⁶ using a simple one step reduction and simultaneous acetylation process achieved by adding thioacetic acid to a pyridine solution containing ManNAc6N₃.⁴⁷ Neu5Ac9NAc was readily obtained from ManNAc6NAc and pyruvate with a 78% yield using a reaction catalyzed by a sialic acid aldolase from *Pasteurella multocida* (PmAldolase).⁴⁵

One-pot Multienzyme (OPME) Synthetic Approach for Facile Production of Neu5Ac9NAc α 3Gal β 4Glc β -ProNH₂ for Glycan Microarray Studies. ManNAc6NAc was also used directly for synthesizing a sialoside analog Neu5Ac9NAc α 3Gal β 4Glc β ProN₃ via an efficient one-pot multienzyme (OPME) sialylation system^{46,48,49} containing PmAldolase,⁴⁵ a CMP-sialic acid synthetase from *Neisseria meningitidis* (NmCSS),⁴⁴ and *Pasteurella multocida* sialyltransferase 1 M144D mutant (PmST1M144D).⁵⁰ Neu5Ac9NAc α 3Gal β 4Glc β ProN₃ was obtained in 84% yield and was readily converted in a quantitative yield to Neu5Ac9NAc α 3Gal β 4Glc β -ProNH₂ by catalytic hydrogenation using H₂ in the presence of Pd/C.⁵¹

Glycan Microarray Studies of Viral Proteins and Human Siglec-9 Binding to Sialosides Containing Neu5,9Ac₂ and Neu5Ac9NAc. The synthesized glycan Neu5Ac9NAc α 3Gal β 4Glc β ProNH₂ was added to our previously described glycan library for glycan microarray studies.²⁸ Proteins known to recognize Neu5,9Ac₂ were tested, including immunoglobulin Fc-fused human Siglec-9-Fc (hSiglec-9),⁵² as well as viral proteins (hemagglutinins in an esterase inactivated form) of porcine torovirus (PToV-P4-Fc, PToV) and bovine coronavirus (BCoV-Mebus-Fc, BCoV).¹³ The results were compared to those obtained using Neu5,9Ac₂ α 3Gal β 4Glc β ProNH₂ and Neu5Ac α 3Gal β 4Glc β -ProNH₂ lacking 9-modification at sialic acid as probes on the same slides (Figure 3). Results from glycan microarray studies (Figure 3A) showed that hSiglec-9 prefers binding to Neu5Ac-glycan without 9-modification at sialic acid (white column), but with observable binding to both Neu5Ac9NAc (black column) and Neu5,9Ac₂ (gray column)-containing glycans. In comparison, both PToV and BCoV prefer binding to both sialoglycans with 9-modifications such as Neu5Ac9NAc and Neu5,9Ac₂. BCoV is a known viral protein probe for 9OAc sialic acids (both Neu5,9Ac₂ and Neu5Gc9Ac) and especially 7,9-di-OAc sialic acids.¹³ The more specific 9OAc-sialic acid viral probe PToV (prefers Neu5,9Ac₂ over Neu5Gc9Ac)¹³ binds to both 9OAc and 9NAc glycans although it shows slightly stronger binding to the 9OAc derivative. The specific recognition of 9-




Figure 3. Sialoglycan microarray binding specificity studies of human Siglec-9 (hSiglec-9-Fc, hSiglec-9), porcine torovirus hemagglutinin-esterase (PToV), and bovine coronavirus hemagglutinin-esterase (BCoV; both PToV and BCoV were mutated to ablate their esterase activity) toward Neu5Ac9NAc α 3Gal β 4Glc β ProNH₂ (black columns), Neu5,9Ac₂ α 3Gal β 4Glc β ProNH₂ (gray columns), and Neu5Ac α 3Gal β 4Glc β ProNH₂ (white columns) without esterase treatment (A) or treated with esterase active PToV (B).

modified sialic acid by esterase-inactive PToV and BCoV, the stability of 9NAc, and the esterase lability of 9OAc were evident by comparing the binding study results of esterase active PToV-treated samples (Figure 3B) with those nontreated ones (Figure 3A).

Feeding of Free Neu5Ac9NAc and Neu5,9Ac₂ to Human Cells Followed by Measuring Cell Surface Incorporation with PToV Probes. Human Burkitt lymphoma B (BJA-B K20) cells are hypo-sialylated due to the lack of UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase,⁵³ therefore maximizing exogenous sialic acid incorporation. After 3 days of feeding with either 1 mM of free Neu5Ac9NAc or Neu5,9Ac₂, cells were probed with PToV-P4-Fc (PToV) with or without active esterase to measure cell surface incorporation of the sugars. Both Neu5Ac9NAc and Neu5,9Ac₂ were detected on the cell surface, but only Neu5Ac9NAc-fed cells were resistant to virus-hemagglutinin-esterase activity (Figure 4).

Feeding of Free Neu5Ac, Neu5,9Ac₂, and Neu5Ac9NAc to Human BJA-B K20 Cells Followed by Measuring Cell Surface Incorporation of Sialic Acids with DMB Derivatization and HPLC Analysis. BJA-B K20 cells were fed with 1 mM Neu5Ac (Figure 5A), Neu5,9Ac₂ (Figure 5B), or Neu5Ac9NAc (Figure 5C) for 3 days. The cell membranes were recovered, and sialic acids were released with *Vibrio cholerae* neuraminidase.^{34,54} The released sialic acids were derivatized with DMB and subjected to HPLC analysis which showed incorporation of each sugar on the cell membrane. Although BJA-B K20 cells are hypo-sialylated, they still express




Figure 4. Detection of incorporation of Neu5,9Ac₂ or Neu5Ac9NAc into hypo-sialylated human lymphoma BJA-B K20 cells. Cells were fed (or not) for 3 days with free Sias (1 mM) and then stained with PTOV in inactive (mutated, solid black line) or active (dashed black line) O-acetyltransferase forms. The binding was analyzed by flow cytometry.




Figure 5. DMB derivatization and HPLC analysis of sialic acids released from cell membranes of BJA-B K20 fed for 3 days with 1 mM Neu5Ac (A), Neu5,9Ac₂ (B) or Neu5Ac9NAc (C).

some Neu5Ac, as can be observed on the cell membrane of Neu5,9Ac₂ or Neu5Ac9NAc-fed cells (Figure 5B and C).

Monitoring Turnover of Neu5Ac9NAc and Neu5,9Ac₂ on the Cell Surface. After feeding BJA-B K20 cells for 2 days with either 1 mM of free Neu5Ac9NAc or Neu5,9Ac₂, the sugars were removed from the cell culture medium (day 0), and the presence of the sialic acids on the cell surface was monitored for 4 days with the PTOV (esterase inactive) probe (Figure 6). The turnover of Neu5,9Ac₂ on the cell surface was shown to be faster than that of Neu5Ac9NAc.

Feeding of Neu5Ac, Neu5,9Ac₂, and Neu5Ac9NAc to Human BJA-B K20 Cells Followed by Measuring Cell Surface Incorporation with Human CD22-Fc. BJA-B K20




Figure 6. Monitoring turnover of incorporated Neu5,9Ac₂ (A) and Neu5Ac9NAc (B) in BJA-B K20 cells. Cells were fed for 2 days with 1 mM of Neu5,9Ac₂ or Neu5Ac9NAc. After 2 days, the feeding with Neu5,9Ac₂ or Neu5Ac9NAc was stopped, and the turnover of these sugars was measured with the PTOV (esterase inactive) probe by flow cytometry in the course of 4 days. The expected amount of sugar was calculated on the basis of the initial MFI (Mean Fluorescence Intensity) at day 0 and the following cell doubling.

cells were fed with 3 mM Neu5Ac, Neu5Ac9NAc, or Neu5,9Ac₂. After 3 days of feeding, BJA-B K20 cells were probed with human CD22-Fc/Siglec-2 to detect ligands (Figure 7). It was found that there was a high expression of




Figure 7. Probing Neu5Ac, Neu5,9Ac₂ or Neu5Ac9NAc-fed human lymphoma BJA-B K20 cells with CD22-Fc to detect ligands. Cells were fed for 3 days with 3 mM Neu5Ac (solid black line), Neu5,9Ac₂ (dashed black line), or Neu5Ac9NAc (dotted line) and then stained with human CD22-Fc to detect ligands (and compared with nonfed cells, gray). The binding was analyzed by flow cytometry.

CD22 ligand when the cells were fed with Neu5Ac (Figure 7A). However, when the cells were grown in the presence of Neu5Ac9NAc, the expression of the CD22 ligand was minimal (Figure 7C). The level of CD22 ligand expressed on the cells fed with Neu5,9Ac₂ (Figure 7B) was between the two, which could be explained by possible partial de-O-acetylation of Neu5,9Ac₂. This supports previous observations that human CD22-Fc recognizes α 2-6-linked sialosides with Neu5Ac or N-glycolylneuraminic acid (Neu5Gc) as the preferred naturally existing sialic acid form and 9-O-acetyl modification on sialic acid blocks the binding.^{6,7} In addition, it demonstrates further that Neu5Ac9NAc can be a suitable substituent for natural Neu5,9Ac₂ which has improved stability.

Comparison of Sialosides Containing Neu5Ac9NAc and Neu5,9Ac₂ Using Molecular Dynamics. Classical molecular dynamics (MD) simulations of

Neu5,9Ac₂α3Galβ4GlcβProN₃ and Neu5Ac9NAcα3-Galβ4GlcβProN₃ were carried out to investigate any differences that the chemical modification would introduce into the conformational ensemble in aqueous solution. The simulations were based on the GLYCAM carbohydrate force field⁵⁵ and TIP3P water model,⁵⁶ with added electrostatic parameters for describing the N-acetylation. The total length of the simulation exceeded 2 μs for each of the two sialosides studied, which has previously been noted to provide good sampling of oligosaccharide conformational degrees of freedom.⁵⁷

Figure 8 shows the results of our simulations projected onto the (ϕ , ψ) dihedral angles of the glycosidic linkage between Sia




Figure 8. Free energy diagram of glycosidic linkage in solution for Neu5Ac9NAcα3Galβ4GlcβProN₃ and the corresponding Neu5,9Ac₂α3Galβ4GlcβProN₃. The 9NAc and 9OAc analogs are compared in the left and right panels. The predominant states are almost identical for the two molecules, indicating that the 9NAc and 9OAc sialosides have similar structures in solution. The 9OAc sialoside has a sparsely populated state not seen in 9NAc (dotted box).

and Gal. The main observation is that the conformational ensembles of the N-acetylated and O-acetylated sialosides are highly equivalent. There are three free energy basins on the positive ψ half of the plot with the same shape, and the relative free energies between the basins are similar to within 1 kcal/mol.

The classical MD simulation approximates the effect of the chemical modification as changing the force field parameters at the acetylation site. In particular, only the partial charges on the N-acetyl functional group are modified. To test this assumption, we used density functional theory to calculate the difference in electrostatic potential as a result of changing O-acetylation to N-acetylation.⁵⁸ The results are shown in Figure 9, which confirm that the change in electrostatic potential is entirely localized to the acetylation site, which lends credence to the results of the classical simulations.

The O-acetylated sialoside appears to display a slightly higher amount of flexibility, as there exists a cluster of conformations at 3–4 kcal/mol higher in energy than the minimum (dotted box in Figure 8) where ψ adopts negative values. These conformations represent far less than 1% of the whole ensemble but are nonetheless interesting to investigate for understanding the effects of chemical modification. The overall conformational ensemble is shown in the top panel of Figure 10, and the negative- ψ conformations are shown in the bottom panel, indicating that the position and orientation of Sia differ significantly between the full set and subset of conformations. The sialoside spends most of its time in an extended conformation, but occasionally adopts a hairpin-like structure where the C7–C9 part of Sia bends back toward Glc. These hairpin conformations were not observed in the N-acetylated




Figure 9. Top: Calculated electrostatic potential for a representative sialoside structure extracted from simulations of Neu5Ac9NAcα3-Galβ4GlcβProN₃ using the B3LYP/6-31G* level of theory^{60,61} and plotted on the $\rho = 0.002$ density isosurface. Bottom: Electrostatic potential difference computed between Neu5Ac9NAcα3Galβ4GlcβProN₃ and Neu5,9Ac₂α3Galβ4GlcβProN₃ plotted on the same isosurface; the O-acetylated potential was calculated by replacing NH with O in the N-acetylated structure. The difference between the potentials is very small, although the NH group has a slightly positive potential relative to O.

Full ensemble of OAc




Figure 10. Superimposed conformations of the 9OAc sialoside simulations, showing only carbohydrate residues and aligned using galactose heavy atoms. One representative conformation is shown in licorice representation (C, gray; O, red; N, blue; H, white). Top: The full conformational ensemble of Neu5,9Ac₂α3Galβ4GlcβProN₃ with snapshots taken 10 ns apart (Neu5,9Ac₂, yellow; Gal, green; Glc, blue). Bottom: The conformations corresponding to the dotted box in Figure 8 with snapshots taken 10 ps apart. These rarely seen conformations were only visited in the O-acetylated simulations but are only a small part of the ensemble, being higher in free energy.

ensemble. The sialosides formed few intramolecular hydrogen bonds, which suggests that the chemical modification alters the conformational ensemble *via* the local hydrogen bonding network of the solvent,⁵⁹ a more detailed analysis of solvent degrees of freedom is needed to test this hypothesis.

DISCUSSION

This study began with the realization that even with the best of precautions, some de-*O*-acetylation occurs during preparation of sialoglycan arrays. In retrospect, definite conclusions to date by us and by others about 9-*O*-acetylation on such arrays can only be made in instances where binding is exclusively to the *O*-acetylated sialoglycan spot, and not to the corresponding non-*O*-acetylated one.¹³ And even in this instance the ideal conclusive proof is to show loss of binding upon pretreatment with a Nidovirus 9-*O*-acetyl esterase (such as PToV and BCoV).¹³ To be clear, this does not overtly contradict our previous studies^{16–19,27–32} as we rarely commented about the role of *O*-acetylation except when selective binding occurred.¹³ But it is important to prevent inadvertent misinterpretation of some of our published raw data such as binding patterns and heat maps, by readers who may not know the subtleties and to alert others who may be studying these fascinating acetyl esters that modify important members of the vertebrate sialome.

Here, we have presented a very practical solution to long-standing technical problems that have markedly inhibited progress in studies of the biology of 9-*O*-acetylated sialic acids. The strategy is replacing the labile *OAc* group with a more stable *NAC* group, which otherwise behaved functionally in a generally similar manner on microarrays, incorporation into cells after feeding, and subsequent expression on cell surfaces. The single obvious difference in both the array and cell feeding studies with or without binding studies using human CD22-Fc as a probe was the general resistance of the *NAC* group to both viral esterases and to endogenous cellular esterases. Of course in both instances we cannot rule out a very low level of esterase cleavage, and further studies are required.

The lack of cell lines that consistently express 9*OAc* sialic acids has also been a great limitation with cellular studies. The incorporation of the 9*NAC* molecules by feeding now allows one to ask if the cells become targets for infection by viruses that normally recognize the 9*OAc* group. Many other previously well-recognized but poorly studied biological or pathological roles of sialic acid *O*-acetylation may now become accessible. For example, while the disialoganglioside GD3 appears to protect cells from apoptosis, the unstable 9*OAcGD3* is said to enhance apoptosis.^{20–24} If feeding of cells expressing GD3 results in expression of 9*NACGD3*, this phenomenon can now be better studied. Likewise the potential role of *O*-acetylation in regulating Siglec functions could also now be better explored.

The rapid turnover of the *O*-acetyl groups in fed cells is very likely due to the action of a cytosolic sialic acid esterase we previously described.^{42,62–64} We predict the *NAC*-form to be resistant to this enzyme. The slower but continued turnover of the *N*-acetyl form is likely due to continued dilution by cell doubling. Assuming that the Neu5Ac9*NAC* is first degraded into Man*NAC6NAC*, the question also arises as to whether this molecule can be further metabolized. Biochemical analysis can be used to track possible outcomes.

In principle, the same approach could be tried to address difficulties in the studies of 4-*O*-acetyl sialic acid and even more labile esters known to occur naturally at the 7- or 8-positions of

sialic acids.² A paper recently appeared showing an example of how this approach can work for 4-*O*-acetyl group.⁶⁵ The approach could also potentially address the strong propensity of *O*-acetyl groups to migrate from one position to another.^{2,54,66,67} Even at near neutral/physiological pH values under which no significant de-*O*-acetylation occurred, the 7-*O*-acetyl group of Neu5,7Ac₂ can easily migrate to the 9-position, producing Neu5,9Ac₂ (possibly via an 8-position intermediate). Even more remarkably, the di-*O*-acetyl form Neu5,7,9Ac₃ quickly yields a dynamic equilibrium mixture of Neu5,7,9Ac₃ and Neu5,8,9Ac₃, in a molar ratio of ~1:1. Both the di-*O*-acetyl forms and the transient mono-8-*O*-acetyl form have been virtually impossible to study. While even more challenging to synthesize, *N*-acetyl versions of the 7,9-di-*O*-acetyl, 8,9-di-*O*-acetyl, 4,7-di-*O*-acetyl, and 4,9-di-*O*-acetyl molecules that are known to occur in nature, as well as the 7,8,9-tri-*O*-acetyl entity, could be considered. Preparation of the *NAC* versions of these molecules could give the first hope of potentially understanding functions of the evanescent 8-*O*-acetyl group, and of differentiating between the recognition functions of 7,9- and 8,9-di-*O*-acetyl sialic acids, which are normally in a dynamic equilibrium state. Nevertheless, it remains to be seen if *NAC* analogs of such molecules recapitulate the conformation and biology of their natural counterparts, and also if they can all be efficiently incorporated by feeding the cells.

Conclusions. We present an efficient one-pot multienzyme method to synthesize a sialoglycan containing Neu5Ac9*NAC* and demonstrate that substituting the C9-oxygen atom in Neu5,9Ac₂ with a nitrogen atom is an effective approach to study sialosides containing unstable 9-*O*-acetyl sialic acid. Computational molecular dynamics simulations, mammalian cell feeding with or without binding studies using human CD22-Fc as a probe, and glycan microarray studies showed that Neu5Ac9*NAC*-sialoglycan structurally and conformationally resembles Neu5,9Ac₂-sialoglycan but with much improved stability in handling and study. This simple approach therefore opens up numerous research opportunities to gain a new understanding of the biology that depends on the modification of sialic acids with an *O*-acetyl group at various positions that have heretofore proved intractable.

METHODS


Materials. Cy3 and PE affinitive goat antihuman IgG (H+L) antibodies were obtained from Jackson ImmunoResearch Laboratories. *Arthrobacter ureafaciens* sialidase (AUS) and *Vibrio cholerae* neuraminidase were purchased from EY Laboratories and Roche, respectfully. Chemical and biological material were purchased from commercial sources and used as received. Nuclear magnetic resonance (NMR) spectra were recorded in the NMR facility of the University of California, Davis on a Bruker Avance-400 NMR spectrometer (400 MHz for ¹H, 100 MHz for ¹³C). Chemical shifts are reported in parts per million (ppm) on the δ scale. High resolution electrospray ionization (ESI) mass spectra were obtained using a Thermo Electron LTQ-Orbitrap Hybrid MS at the Mass Spectrometry Facility at the University of California, Davis. Column chromatography was performed using RediSep Rf silica columns or an ODS-SM column (51 g, 50 μ m, 120 Å, Yamazen) on the CombiFlash Rf 200i system. Thin layer chromatography (TLC) was performed on silica gel plates (Sorbert Technologies) using anisaldehyde sugar stain for detection. Gel filtration chromatography was performed with a column (100 cm \times 2.5 cm) packed with Bio-Gel P-2 Fine resins (Bio-Rad). *Pasteurella multocida* sialic acid aldolase (PmNanA), *Neisseria meningitidis* CMP-sialic acid synthetase (NmCSS), and *Pasteurella multocida* sialyltransferase 1 M144D mutant (PmST1M144D) were expressed and purified as described previously.

Preparation of Hemagglutinin-Esterase Probes for 9-OAc Sia. The sequences for expressing the ectodomains of the hemagglutinin-esterase (HE) of two nidoviruses (PTov-P4 and BCoV-Mebus) were synthesized by GenScript (Piscataway, NJ), with varied degrees of insect codon optimization. To generate the probe molecules, the esterase domains were inactivated by changing the active site of the Ser residue to Ala by site-directed mutagenesis using Q5 mutagenesis (New England Biolabs). The HE proteins were linked to the baculovirus gp64 signal sequence peptide at their N-termini and the C-terminus fused to a linker containing a thrombin cleavage sequence, the Fc domain of Human IgG1, and a 6-His sequence. Constructs were cloned into pFastBac-1 (Life Technologies) to generate recombinant bacmids following the manufacturer's protocol. Recombinant baculoviruses were recovered by transfection of the bacmids into Sf9 insect cells using Cellfectin II (Life Technologies). Viruses were used to infect suspension High Five cells and the supernatant harvested 2–3 days postinfection. The proteins were purified by binding to a HiTrap ProteinG HP 5 mL column (GE Healthcare Life Sciences, Piscataway, NJ) and eluted with 0.1 M citrate, at pH 3.0 (pH neutralization to 7.8 with 1 M of Tris-HCl, pH 9.0), using the ÄKTA FPLC system (GE Healthcare Life Sciences). The HE-Fc containing fractions were dialyzed in PBS and concentrated using 30 kDa Amicon Ultra-15 filters (EMD Millipore). Purified proteins were stored at -80°C in aliquots.

Cell Culture and Sugar Supplementation. BJA-B K20 cells were propagated as a suspension with RPMI 1640 medium and supplemented with 10% heat-inactivated fetal calf serum, 2 mM glutamine, 100 U of penicillin per mL, and 100 μg of streptomycin per mL in a humidified 5% CO_2 , 37°C atmosphere as described previously.⁵³ For medium supplementation, Neu5Ac9NAc and Neu5,9Ac₂ were dissolved in PBS, titrated to a neutral pH, and filter sterilized. The sugars were added at the indicated concentrations. For the sugar turnover experiments, the cells were fed as mentioned above and on day 0, the cell culture medium was switched to RPMI 1640 medium, supplemented with 1% Nutridoma (Roche), 100 U of penicillin per mL, and 100 μg of streptomycin without Neu5Ac9NAc and Neu5,9Ac₂.

Flow Cytometry. Cells were washed with PBS and incubated with 10 $\mu\text{g}/\text{mL}$ of PTov probes in PBS with 1% BSA and 10 mM EDTA for 30 min on ice. Subsequently, cells were washed with PBS and incubated with PE goat antihuman IgG antibody for 30 min on ice. After an additional washing step, the cells were analyzed by FACSCalibur (BD).

Synthesis of 2,6-Diacetamido-2,6-dideoxy-D-mannopyranose (ManNAc6NAc) from 2-Acetamido-6-azido-2,6-dideoxy-D-mannopyranose (ManNAc6N₃). To a solution of ManNAc6N₃ (300 mg, 1.14 mmol) in pyridine (8 mL), thioacetic acid (2 mL) was added, and the mixture was stirred at RT for 24 h. The solvent was concentrated *in vacuo*. The crude product was purified by column chromatography (ethyl acetate/methanol = 3:1) to produce ManNAc6NAc (262 mg, 82%) as a colorless amorphous solid. ¹H NMR (400 MHz, D₂O) α -isomer: δ 5.10 (d, J = 1.5 Hz, 1H), 4.31 (dd, J = 4.7, 1.5 Hz, 1H), 4.04 (dd, J = 9.8, 4.7 Hz, 1H), 3.96–3.86 (m, 1H), 3.68–3.33 (m, 3H), 2.07 (s, 3H), 2.03 (s, 3H). β -isomer: δ 5.00 (d, J = 1.7 Hz, 1H), 4.46 (dd, J = 4.5, 1.7 Hz, 1H), 3.82 (dd, J = 9.0, 4.5 Hz, 1H), 3.68–3.33 (m, 4H), 2.11 (s, 3H), 2.04 (s, 3H). ¹³C NMR (100 MHz, D₂O): δ 175.70, 174.77, 174.54, 174.53, 93.01, 74.43, 71.73, 70.24, 68.58, 68.49, 68.20, 53.98, 53.21, 40.27, 40.16, 22.04, 21.90, 21.77. HRMS (ESI) Anal. Calcd for C₁₀H₁₉N₂O₂ [M + H]⁺: 263.1243. Found: 263.1241.




Synthesis of 5,9-Diacetamido-3,5,9-Trideoxy-D-glycero-D-galacto-2-nonulopyranosylonic Acid (Neu5Ac9NAc). To a solution (10 mL) containing ManNAc6NAc (50 mg, 0.19 mmol), sodium pyruvate (210 mg, 1.9 mmol), and Tris-HCl buffer (100 mM, pH 7.5), PmAldolase (3.0 mg) was added, and the reaction was incubated in an isotherm incubator for 48 h at 37°C with agitation at

100 rpm. The reaction was quenched by adding the same volume of ice-cold ethanol and incubating at 4°C for 1 h. The solvent was removed, and the crude product was purified by column chromatography (ethyl acetate/methanol/water = 4:2:1) and followed by a short BioGel P-2 gel filtration column to give Neu5Ac9NAc (55.4 mg, 78%) as a colorless amorphous solid. ¹H NMR (400 MHz, D₂O): δ 4.06–3.92 (m, 3H, H-4, H-6, H-5), 3.80 (ddd, J = 9.0, 7.7, 3.1 Hz, 1H, H-8), 3.59 (dd, J = 14.1, 3.1 Hz, 1H, H-9), 3.44 (dd, J = 9.0, 1.1 Hz, 1H, H-7), 3.24 (dd, J = 14.1, 7.6 Hz, 1H, H-9), 2.22 (dd, J = 12.9, 4.8 Hz, 1H, H-3e), 2.06 (s, 3H, COCH₃), 2.02 (s, 3H, COCH₃), 1.83 (dd, J = 12.9, 11.4 Hz, 1H, H-3a). ¹³C NMR (100 MHz, D₂O): δ 176.64, 174.68, 174.56, 96.36, 70.09, 69.77, 68.77, 67.25, 52.26, 42.76, 39.35, 22.08, 21.82. HRMS (ESI) Anal. Calcd for C₁₃H₂₁N₂O₉ [M – H][–]: 349.1247. Found: 349.1255.

Synthesis of 3-Azidopropyl O-(5,9-Diacetamido-3,5,9-trideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 3)-O- β -D-galactopyranosyl-(1 \rightarrow 4)- β -D-glucopyranoside (Neu5Ac9NAc α 3Gal β 4Glc β ProN₃). Neu5Ac9NAc α 3Gal β 4Glc β ProN₃ was synthesized following the general procedure for enzymatic synthesis of sialosides. Gal β 4Glc β ProN₃⁴⁶ (46 mg, 0.11 mmol), ManNAc6NAc (43 mg, 0.16 mmol), sodium pyruvate (91 mg, 0.83 mmol), and CTP (87 mg, 0.16 mmol) were dissolved in Tris-HCl buffer (100 mM, pH 8.5, 10 mL) containing 20 mM MgCl₂. The pH of the solution was further adjusted to 8.5 with 4 M NaOH. PmAldolase (1.5 mg), NmCSS (2.5 mg), and PmST1M144D (2.5 mg) were added. The reaction was incubated in an isotherm incubator for 24 h at 37°C with agitation at 100 rpm. The reaction was quenched by adding the same volume of ice-cold ethanol and incubating at 4°C for 1 h. The formed precipitates were removed by centrifugation, and the supernatant was concentrated. The residue was purified by passing it through a BioGel P-2 gel filtration column followed by a C18 column (H₂O/CH₃CN = 10:1; 70.9 mg, Yield 84%). ¹H NMR (400 MHz, D₂O): δ 4.52 (dd, J = 12.8, 7.9 Hz, 2H), 4.10 (dd, J = 9.9, 3.2 Hz, 1H), 4.06–3.86 (m, 4H), 3.88–3.53 (m, 13H), 3.54–3.43 (m, 3H), 3.37–3.23 (m, 2H), 2.76 (dd, J = 12.4, 4.6 Hz, 1H), 2.04 (s, 3H), 2.04 (s, 3H), 1.92 (p, J = 6.6 Hz, 2H), 1.81 (t, J = 12.1 Hz, 1H). ¹³C NMR (100 MHz, D₂O): δ 174.92, 174.39, 173.81, 102.60, 102.11, 99.87, 78.17, 75.53, 75.14, 74.78, 74.33, 72.79, 72.73, 69.96, 69.58, 69.36, 68.30, 67.46, 67.34, 61.01, 60.01, 51.66, 47.86, 42.11, 39.61, 28.22, 22.02, 21.81. HRMS (ESI) Anal. Calcd for C₂₈H₄₆N₃O₁₉ [M – H][–]: 756.2787. Found: 756.2792.

Synthesis of 3-Aminopropyl O-(5,9-Diacetamido-3,5,9-trideoxy-D-glycero- α -D-galacto-2-nonulopyranosylonic acid)-(2 \rightarrow 3)-O- β -D-galactopyranosyl-(1 \rightarrow 4)- β -D-glucopyranoside (Neu5Ac9NAc α 3Gal β 4Glc β ProNH₂). A catalytic amount of 10% palladium on charcoal (Pd/C) was added to the solution of Neu5Ac9NAc α 3Gal β 4Glc β ProN₃ (5.6 mg) in H₂O (1 mL). The mixture was stirred under a hydrogen atmosphere for 3 h. The solution was diluted with MeOH (2 mL) and passed through a filter to remove the catalyst. The solvent was concentrated *in vacuo* to give Neu5Ac9NAc α 3Gal β 4Glc β ProNH₂ (5.4 mg, quant.) as a colorless amorphous solid. HRMS (ESI) Anal. Calcd for C₂₈H₄₈N₃O₁₉ [M – H][–]: 730.2882. Found: 730.2914.



Reduction of Neu5,9Ac₂α3Galβ4GlcβProN₃ to Neu5,9Ac₂α3Galβ4GlcβProNH₂. Neu5,9Ac₂α3Galβ4GlcβProN₃ (20 mg) was dissolved in H₂O/MeOH (3 mL, 2:1 by volume), and a drop of glacial acetic acid was added. The mixture was stirred under a hydrogen atmosphere in the presence of 10% palladium on charcoal for 1 h. The reaction mixture was then passed through a HyperSep C18 cartridge (1 g, 40–60 μm , Thermo) and eluted with water. The collection fraction containing the desired product was collected and

lyophilized. The obtained white powder was stored at $-20\text{ }^{\circ}\text{C}$ for long-term storage.

HPLC Analysis of Neu5,9Ac₂α3Galβ4GlcβProNH₂. Neu5,9Ac₂α3Galβ4GlcβProNH₂ was first hydrolyzed in glacial acetic acid (2 M) for 3 h at 80 °C. AUS hydrolysis was performed on the basis of the company protocol. DMB derivatization was performed as was reported previously.⁶⁸ The DMB-derivatized samples were analyzed on a Dionex Ultra3000 HPLC System using a Phenomenex Gemini 5 μm C18 250 × 4.6 mm HPLC column at RT. The fluorescence was detected at 448 nm using excitation at 373 nm.

Glycan Microarray Screening. Glycan microarrays were fabricated using epoxide-derivatized slides (Corning by Thermo Fisher Scientific) and Arrayit SpotBot Extreme Microarray Spotter as previously described.^{17,28,53} The arrays were printed with Stealth SMP3 microarray spotting pins from ArrayIt (Sunnyvale, CA, USA) generating 100 μm diameter spots. Four pins were used, with each pin printing four replicate spots/well. The glycoconjugates were prepared at 100 μM concentration in an optimized printing buffer (300 mM phosphate buffer, pH 8.4). They were then distributed into a 384-well source plate in 20 μL per well. To monitor printing quality, four replicate wells of human IgG (Jackson ImmunoResearch) was used at 100 μg/mL (in PBS) for each printing pin. One complete array was printed on each slide (within approx 12 h/~28 slides). The humidity level in the arraying chamber was maintained at about 60–65% during printing. Printed slides were left on a spotter deck overnight, allowing humidity to drop to ambient levels. Printed glycan microarray slides were blocked by 50 °C prewarmed blocking solution (0.1 M Tris-HCl, 0.05 M ethanolamine, pH 9.0, 1 h). They were then washed twice with 50 °C prewarmed water and dried. Slides were packed, vacuum-sealed, and stored at RT until used.

To ensure the quality of the printed slides, we used one or two slides for quality control using antibodies, lectins, and serum samples with known glycan binding specificity. For glycan array binding, slides were fitted in a multiwell microarray hybridization cassette (AHC4 × 8S, ArrayIt, Sunnyvale, CA, USA) to divide into eight subarrays. The subarrays were blocked with Ovalbumin (1% w/v) in PBS (pH 7.4) for 1 h at RT, with gentle shaking. For the esterase activity studies on the array, after removing the blocking solution, slides were first treated with 20 μg/mL PToV-P4-Fc (esterase active) in blocking buffer for 2 h at RT and then washed with PBS in 0.1% Tween 20 and then PBS (10 min/wash with shaking). Subsequently, diluted protein samples in blocking solution with various concentrations were added to each subarray (both esterase treated and nontreated slides). After incubating the protein samples for 2 h at RT with gentle shaking, the slides were washed with PBS in 0.1% Tween 20 and then PBS (10 min/wash with shaking). Cy3 affinitive goat antihuman IgG (H+L) antibody (Jackson ImmunoResearch) was used for the detection of BCoV-Mebus-Fc, PToV-P4-Fc, and human Siglec-9-Fc. Diluted antibody in PBS was added to the subarrays and incubated for 1 h at RT. They were washed with PBS in 0.1% Tween 20, PBS, and water (10 min/wash with shaking) and dried. The microarray slides were scanned using a Genepix 4000B microarray scanner (Molecular Devices Corp., Union City, CA, USA) at 100% laser power, PMT Gain 450, and 10 μM pixels. Data analysis was performed using Genepix Pro 7.0 analysis software (Molecular Devices Corp., Union City, CA), and the outputs were saved as gpr and jpg files. The gpr files then were saved as xls. The data were further analyzed with Excel. Local background subtraction was performed, and data were plotted separately for each subarray. The binding specificity to glycoconjugates for each protein was plotted based on the RFU (Relative Fluorescence Units), average fluorescence value for four replicates, versus glycan IDs. The standard deviations were calculated, and the error bars were found for each glycan binding. The final graphs (Figure 3) were plotted based on the tested proteins versus RFU for each glycan in GraphPad Prism 5.

Comparison of N-Acetylated and O-Acetylated Sialosides Using Molecular Dynamics. The molecular dynamics simulations were carried out using the AMBER15 software suite, principally using *tLeap* for setup and *pmemd.cuda* for dynamics.⁶⁹ The simulations employed the GLYCAM force field and TIP3P water model; the

sialoside was solvated in a rectangular water box with roughly 1000 water molecules. The simulations used a 2.0 fs time step and a Langevin thermostat set to 298.15 K and a collision frequency of 1.0 ps⁻¹. The equilibration simulations used a Berendsen barostat set to 1.0 atm, a compressibility of $44.6 \times 10^{-6} \text{ bar}^{-1}$, and a time constant of 1.0 ps⁻¹. The production simulations were carried out at constant volume. The particle mesh Ewald method was used to treat long-range electrostatics with a real-space cutoff of 8.0 Å. All input files and parameters are provided as part of the Supporting Information.

The parameters for the N-acetyl functional group were developed following the procedure outlined in ref 70. The initial guess force field for Neu5Ac9NAc was constructed by replacing the 9-O-acetyl group with a copy of the 5-N-acetyl group and the corresponding force field parameters. After 100 ps of equilibration in the NPT ensemble, we generated 50 ns of NVT dynamics and saved 50 snapshots at 1 ns intervals. These snapshots were energy-minimized in the TeraChem software package with the dihedral angles constrained, employing the restricted Hartree–Fock method and 6-31G* basis set.^{71–74} The energy-minimized geometries were used as input to a restrained electrostatic potential calculation carried out using the R.E.D. server using the RESP-C2 charge model as appropriate for the GLYCAM force field.⁵⁵ In the charge optimization, the charges for the terminal hydroxyl group were kept at the initial values and all hydrogen charges set to zero. The resulting final set of charges on the N-acetyl group has a slightly different total charge from the original O-acetyl group; the charge on the carbon atom that the N-acetyl group is bonded to is increased by +0.0014, instead of the +0.008 value used for O-acetyl. An analogous calculation was carried out for the ProN₃ functional group to obtain a complete set of charges for Neu5,9Ac₂α3Galβ4GlcβProN₃. The valence force field parameters for ProN₃ were copied over from the general AMBER force field (GAFF)⁷⁵ by comparing GLYCAM atom types to GAFF atom types.

After the parametrization was finished, the production simulations of Neu5,9Ac₂α3Galβ4GlcβProN₃ and Neu5Ac9NAcα3Galβ4GlcβProN₃ were carried out to collect data for the free energy plots. The simulations were unbiased and each simulation run for 2.5 μs, and configurations were saved every 10 ps. The free energy plot was constructed by projecting the ensemble of configurations onto the selected dihedral degrees of freedom.

■ ASSOCIATED CONTENT

📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acchembio.6b00928.

NMR spectra for products as well and computational residue files, parameter file, and input files for molecular dynamics simulations (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

*Tel.: +1 858-534-2214. Fax: +1 858-534-5611. E-mail: alvarki@ucsd.edu.

*Tel.: +1 530-754-6037. Fax: +1 530-752-8995. E-mail: xiichen@ucdavis.edu.

ORCID

Lee-Ping Wang: 0000-0003-3072-9946

Xi Chen: 0000-0002-3160-614X

Author Contributions

||These authors contributed equally.

Notes

The authors declare the following competing financial interest(s): H.Y. and X.C. are co-founders of Glycohub, Inc., a company focused on the development of carbohydrate based reagents, diagnostics, and therapeutics. Glycohub, Inc. played

no role in the design, execution, interpretation, or publication of this study.

ACKNOWLEDGMENTS

This work was supported by United States National Institutes of Health grants R01HD065122 (to X.C.) and R01GM32373 (to A.V.).

ABBREVIATIONS

AUS, *Arthrobacter ureafaciens* sialidase; DMB, 1,2-diamino-4,5-methylenedioxybenzene; ESI, electrospray ionization; HE, hemagglutinin-esterase; HPLC, high performance liquid chromatography; HRMS, high resolution mass spectrometry; ManNAc6NAc, 6-acetamido-6-deoxy-N-acetylmannosamine; MD, molecular dynamics; MS, mass spectrometry; Neu5Ac, N-acetylneuraminic acid; Neu5,9Ac₂, 9-O-acetyl-N-acetylneuraminic acid; Neu5Ac9NAc, 9-acetamido-9-deoxy-N-acetylneuraminic acid; NMR, nuclear magnetic resonance; OPME, one-pot multienzyme; Sia, sialic acid; TLC, thin layer chromatography

REFERENCES

(1) Lundblad, A. (2015) Gunnar Blix and his discovery of sialic acids. Fascinating molecules in glycobiology. *Uppsala J. Med. Sci.* 120, 104–112.

(2) Schauer, R. (1982) Chemistry, metabolism, and biological functions of sialic acids. *Adv. Carbohydr. Chem. Biochem.* 40, 131–234.

(3) Rogers, G. N., Herrler, G., Paulson, J. C., and Klenk, H. D. (1986) Influenza C virus uses 9-O-acetyl-N-acetylneuraminic acid as a high affinity receptor determinant for attachment to cells. *J. Biol. Chem.* 261, 5947–5951.

(4) Varki, A., and Kornfeld, S. (1980) An autosomal dominant gene regulates the extent of 9-O-acetylation of murine erythrocyte sialic acids. A probable explanation for the variation in capacity to activate the human alternate complement pathway. *J. Exp. Med.* 152, 532–544.

(5) Shi, W. X., Chammas, R., Varki, N. M., Powell, L., and Varki, A. (1996) Sialic acid 9-O-acetylation on murine erythrocyte cells affects complement activation, binding to I-type lectins, and tissue homing. *J. Biol. Chem.* 271, 31526–31532.

(6) Sjoberg, E. R., Powell, L. D., Klein, A., and Varki, A. (1994) Natural ligands of the B cell adhesion molecule CD22 beta can be masked by 9-O-acetylation of sialic acids. *J. Cell Biol.* 126, 549–562.

(7) Kelm, S., Schauer, R., Manuguerra, J. C., Gross, H. J., and Crocker, P. R. (1994) Modifications of cell surface sialic acids modulate cell adhesion mediated by sialoadhesin and CD22. *Glycoconjugate J.* 11, 576–585.

(8) Herrler, G., Rott, R., Klenk, H. D., Muller, H. P., Shukla, A. K., and Schauer, R. (1985) The receptor-destroying enzyme of influenza C virus is neuraminidase-O-acetyltransferase. *EMBO J.* 4, 1503–1506.

(9) Herrler, G., Reuter, G., Rott, R., Klenk, H. D., and Schauer, R. (1987) N-acetyl-9-O-acetylneuraminic acid, the receptor determinant for influenza C virus, is a differentiation marker on chicken erythrocytes. *Biol. Chem. Hoppe-Seyler* 368, 451–454.

(10) Muchmore, E. A., and Varki, A. (1987) Selective inactivation of influenza C esterase: a probe for detecting 9-O-acetylated sialic acids. *Science* 236, 1293–1295.

(11) Enard, W., Khaitovich, P., Klose, J., Zollner, S., Heissig, F., Giavalisco, P., Nieselt-Struwe, K., Muchmore, E., Varki, A., Ravid, R., Doxiadis, G. M., Bontrop, R. E., and Paabo, S. (2002) Intra- and interspecific variation in primate gene expression patterns. *Science* 296, 340–343.

(12) Schauer, R. (1978) Characterization of sialic acids. *Methods Enzymol.* 50, 64–89.

(13) Langereis, M. A., Bakkers, M. J., Deng, L., Padler-Karavani, V., Vervoort, S. J., Hulswit, R. J., van Vliet, A. L., Gerwig, G. J., de Poot, S. A., Boot, W., van Ederen, A. M., Heesters, B. A., van der Loos, C. M.,

van Kuppeveld, F. J., Yu, H., Huizinga, E. G., Chen, X., Varki, A., Kamerling, J. P., and de Groot, R. J. (2015) Complexity and diversity of the mammalian sialome revealed by nidovirus virolectins. *Cell Rep.* 11, 1966–1978.

(14) Song, H., Qi, J., Khedri, Z., Diaz, S., Yu, H., Chen, X., Varki, A., Shi, Y., and Gao, G. F. (2016) An Open Receptor-Binding Cavity of Hemagglutinin-Esterase-Fusion Glycoprotein from Newly-Identified Influenza D Virus: Basis for Its Broad Cell Tropism. *PLoS Pathog.* 12, e1005411.

(15) Cheresch, D. A., Reisfeld, R. A., and Varki, A. P. (1984) O-Acetylation of disialoganglioside GD3 by human melanoma cells creates a unique antigenic determinant. *Science* 225, 844–846.

(16) Padler-Karavani, V., Tremoulet, A. H., Yu, H., Chen, X., Burns, J. C., and Varki, A. (2013) A simple method for assessment of human anti-Neu5Gc antibodies applied to Kawasaki disease. *PLoS One* 8, e58443.

(17) Padler-Karavani, V., Song, X., Yu, H., Hurtado-Ziola, N., Huang, S., Muthana, S., Chokhawala, H. A., Cheng, J., Verhagen, A., Langereis, M. A., Kleene, R., Schachner, M., de Groot, R. J., Lasanajak, Y., Matsuda, H., Schwab, R., Chen, X., Smith, D. F., Cummings, R. D., and Varki, A. (2012) Cross-comparison of protein recognition of sialic acid diversity on two novel sialoglycan microarrays. *J. Biol. Chem.* 287, 22593–22608.

(18) Lu, Q., Padler-Karavani, V., Yu, H., Chen, X., Wu, S. L., Varki, A., and Hancock, W. S. (2012) LC-MS analysis of polyclonal human anti-Neu5Gc xeno-autoantibodies immunoglobulin G Subclass and partial sequence using multistep intravenous immunoglobulin affinity purification and multienzymatic digestion. *Anal. Chem.* 84, 2761–2768.

(19) Padler-Karavani, V., Hurtado-Ziola, N., Chang, Y. C., Sonnenburg, J. L., Ronaghy, A., Yu, H., Verhagen, A., Nizet, V., Chen, X., Varki, N., Varki, A., and Angata, T. (2014) Rapid evolution of binding specificities and expression patterns of inhibitory CD33-related Siglecs in primates. *FASEB J.* 28, 1280–1293.

(20) Birks, S. M., Danquah, J. O., King, L., Vlasak, R., Gorecki, D. C., and Pilkington, G. J. (2011) Targeting the GD3 acetylation pathway selectively induces apoptosis in glioblastoma. *Neuro. Oncol.* 13, 950–960.

(21) Malisan, F., Franchi, L., Tomassini, B., Ventura, N., Condo, I., Rippo, M. R., Rufini, A., Liberati, L., Nachtigall, C., Kniep, B., and Testi, R. (2002) Acetylation suppresses the proapoptotic activity of GD3 ganglioside. *J. Exp. Med.* 196, 1535–1541.

(22) Malisan, F., and Testi, R. (2002) GD3 in cellular ageing and apoptosis. *Exp. Gerontol.* 37, 1273–1282.

(23) Bhunia, A. K., Schwarzmann, G., and Chatterjee, S. (2002) GD3 recruits reactive oxygen species to induce cell proliferation and apoptosis in human aortic smooth muscle cells. *J. Biol. Chem.* 277, 16396–16402.

(24) Chen, H. Y., and Varki, A. (2002) O-Acetylation of GD3: an enigmatic modification regulating apoptosis? *J. Exp. Med.* 196, 1529–1533.

(25) Herrler, G., Gross, H. J., Imhof, A., Brossmer, R., Milks, G., and Paulson, J. C. (1992) A synthetic sialic acid analogue is recognized by influenza C virus as a receptor determinant but is resistant to the receptor-destroying enzyme. *J. Biol. Chem.* 267, 12501–12505.

(26) Feizi, T., Fazio, F., Chai, W., and Wong, C. H. (2003) Carbohydrate microarrays - a new set of technologies at the frontiers of glycomics. *Curr. Opin. Struct. Biol.* 13, 637–645.

(27) Feng, K. H., Gonzalez, G., Deng, L., Yu, H., Tse, V. L., Huang, L., Huang, K., Wasik, B. R., Zhou, B., Wentworth, D. E., Holmes, E. C., Chen, X., Varki, A., Murcia, P. R., and Parrish, C. R. (2015) Equine and canine influenza H3N8 viruses show minimal biological differences despite phylogenetic divergence. *J. Virol.* 89, 6860–6873.

(28) Deng, L., Bensing, B. A., Thamadilok, S., Yu, H., Lau, K., Chen, X., Ruhl, S., Sullam, P. M., and Varki, A. (2014) Oral streptococci utilize a Siglec-like domain of serine-rich repeat adhesins to preferentially target platelet sialoglycans in human blood. *PLoS Pathog.* 10, e1004540.

- (29) Laubli, H., Pearce, O. M., Schwarz, F., Siddiqui, S. S., Deng, L., Stanczak, M. A., Deng, L., Verhagen, A., Secrest, P., Lusk, C., Schwartz, A. G., Varki, N. M., Bui, J. D., and Varki, A. (2014) Engagement of myelomonocytic Siglecs by tumor-associated ligands modulates the innate immune response to cancer. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 14211–14216.
- (30) Deng, L., Song, J., Gao, X., Wang, J., Yu, H., Chen, X., Varki, N., Naito-Matsui, Y., Galan, J. E., and Varki, A. (2014) Host adaptation of a bacterial toxin from the human pathogen *Salmonella Typhi*. *Cell* **159**, 1290–1299.
- (31) Scobie, L., Padler-Karavani, V., Le Bas-Bernardet, S., Crossan, C., Blaha, J., Matouskova, M., Hector, R. D., Cozzi, E., Vanhove, B., Charreau, B., Blanche, G., Bourdais, L., Tallacchini, M., Ribes, J. M., Yu, H., Chen, X., Kracikova, J., Broz, L., Hejnar, J., Vesely, P., Takeuchi, Y., Varki, A., and Soullilou, J. P. (2013) Long-term IgG response to porcine Neu5Gc antigens without transmission of PERV in burn patients treated with porcine skin xenografts. *J. Immunol.* **191**, 2907–2915.
- (32) Padler-Karavani, V., Hurtado-Ziola, N., Pu, M., Yu, H., Huang, S., Muthana, S., Chokhawala, H. A., Cao, H., Secrest, P., Friedmann-Morvinski, D., Singer, O., Ghaderi, D., Verma, I. M., Liu, Y. T., Messer, K., Chen, X., Varki, A., and Schwab, R. (2011) Human xeno-autoantibodies against a non-human sialic acid serve as novel serum biomarkers and immunotherapeutics in cancer. *Cancer Res.* **71**, 3352–3363.
- (33) Chokhawala, H. A., Huang, S., Lau, K., Yu, H., Cheng, J., Thon, V., Hurtado-Ziola, N., Guerrero, J. A., Varki, A., and Chen, X. (2008) Combinatorial chemoenzymatic synthesis and high-throughput screening of sialosides. *ACS Chem. Biol.* **3**, 567–576.
- (34) Chokhawala, H. A., Yu, H., and Chen, X. (2007) High-throughput substrate specificity studies of sialidases by using chemoenzymatically synthesized sialoside libraries. *ChemBioChem* **8**, 194–201.
- (35) Samraj, A. N., Pearce, O. M., Laubli, H., Crittenden, A. N., Bergfeld, A. K., Banda, K., Gregg, C. J., Bingman, A. E., Secrest, P., Diaz, S. L., Varki, N. M., and Varki, A. (2015) A red meat-derived glycan promotes inflammation and cancer progression. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 542–547.
- (36) Corfield, A. P., Williams, A. J., Clamp, J. R., Wagner, S. A., and Mountford, R. A. (1988) Degradation by bacterial enzymes of colonic mucus from normal subjects and patients with inflammatory bowel disease: the role of sialic acid metabolism and the detection of a novel O-acetylsialic acid esterase. *Clin. Sci.* **74**, 71–78.
- (37) Corfield, A. P., Wagner, S. A., O'Donnell, L. J., Durdey, P., Mountford, R. A., and Clamp, J. R. (1993) The roles of enteric bacterial sialidase, sialate O-acetyl esterase and glycosulfatase in the degradation of human colonic mucin. *Glycoconjugate J.* **10**, 72–81.
- (38) Steenbergen, S. M., Jirik, J. L., and Vimr, E. R. (2009) YjhS (NanS) is required for *Escherichia coli* to grow on 9-O-acetylated N-acetylneuraminic acid. *J. Bacteriol.* **191**, 7134–7139.
- (39) Rangarajan, E. S., Ruane, K. M., Proteau, A., Schrag, J. D., Valladares, R., Gonzalez, C. F., Gilbert, M., Yakunin, A. F., and Cygler, M. (2011) Structural and enzymatic characterization of NanS (YjhS), a 9-O-acetyl N-acetylneuraminic acid esterase from *Escherichia coli* O157:H7. *Protein Sci.* **20**, 1208–1219.
- (40) Phansopa, C., Kozak, R. P., Liew, L. P., Frey, A. M., Farmilo, T., Parker, J. L., Kelly, D. J., Emery, R. J., Thomson, R. I., Royle, L., Gardner, R. A., Spencer, D. L., and Stafford, G. P. (2015) Characterization of a sialate-O-acetyltransferase (NanS) from the oral pathogen *Tannerella forsythia* that enhances sialic acid release by NanH, its cognate sialidase. *Biochem. J.* **472**, 157–167.
- (41) Hayes, B. K., and Varki, A. (1989) O-Acetylation and de-O-acetylation of sialic acids. Sialic acid esterases of diverse evolutionary origins have serine active sites and essential arginine residues. *J. Biol. Chem.* **264**, 19443–19448.
- (42) Takematsu, H., Diaz, S., Stoddart, A., Zhang, Y., and Varki, A. (1999) Lysosomal and cytosolic sialic acid 9-O-acetyltransferase activities can be encoded by one gene via differential usage of a signal peptide-encoding exon at the N terminus. *J. Biol. Chem.* **274**, 25623–25631.
- (43) Schauer, R., and Shukla, A. K. (2008) Isolation and properties of two sialate-O-acetyltransferases from horse liver with 4- and 9-O-acetyl specificities. *Glycoconjugate J.* **25**, 625–632.
- (44) Yu, H., Yu, H., Karpel, R., and Chen, X. (2004) Chemoenzymatic synthesis of CMP-sialic acid derivatives by a one-pot two-enzyme system: comparison of substrate flexibility of three microbial CMP-sialic acid synthetases. *Bioorg. Med. Chem.* **12**, 6427–6435.
- (45) Li, Y., Yu, H., Cao, H., Lau, K., Muthana, S., Tiwari, V. K., Son, B., and Chen, X. (2008) *Pasteurella multocida* sialic acid aldolase: a promising biocatalyst. *Appl. Microbiol. Biotechnol.* **79**, 963–970.
- (46) Yu, H., Chokhawala, H., Karpel, R., Yu, H., Wu, B., Zhang, J., Zhang, Y., Jia, Q., and Chen, X. (2005) A multifunctional *Pasteurella multocida* sialyltransferase: a powerful tool for the synthesis of sialoside libraries. *J. Am. Chem. Soc.* **127**, 17618–17619.
- (47) Shanguan, N., Katukojvala, S., Greenberg, R., and Williams, L. J. (2003) The reaction of thio acids with azides: a new mechanism and new synthetic applications. *J. Am. Chem. Soc.* **125**, 7754–7755.
- (48) Yu, H., Chokhawala, H. A., Huang, S., and Chen, X. (2006) One-pot three-enzyme chemoenzymatic approach to the synthesis of sialosides containing natural and non-natural functionalities. *Nat. Protoc.* **1**, 2485–2492.
- (49) Yu, H., Huang, S., Chokhawala, H., Sun, M., Zheng, H., and Chen, X. (2006) Highly efficient chemoenzymatic synthesis of naturally occurring and non-natural alpha-2,6-linked sialosides: a *P. damsela* alpha-2,6-sialyltransferase with extremely flexible donor-substrate specificity. *Angew. Chem., Int. Ed.* **45**, 3938–3944.
- (50) Sugiarto, G., Lau, K., Qu, J., Li, Y., Lim, S., Mu, S., Ames, J. B., Fisher, A. J., and Chen, X. (2012) A sialyltransferase mutant with decreased donor hydrolysis and reduced sialidase activities for directly sialylating Lewis^x. *ACS Chem. Biol.* **7**, 1232–1240.
- (51) Yu, H., Chokhawala, H. A., Varki, A., and Chen, X. (2007) Efficient chemoenzymatic synthesis of biotinylated human serum albumin-sialoglycoside conjugates containing O-acetylated sialic acids. *Org. Biomol. Chem.* **5**, 2458–2463.
- (52) Zhang, J. Q., Nicoll, G., Jones, C., and Crocker, P. R. (2000) Siglec-9, a novel sialic acid binding member of the immunoglobulin superfamily expressed broadly on human blood leukocytes. *J. Biol. Chem.* **275**, 22121–22126.
- (53) Hinderlich, S., Berger, M., Keppler, O. T., Pawlita, M., and Reutter, W. (2001) Biosynthesis of N-acetylneuraminic acid in cells lacking UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase. *Biol. Chem.* **382**, 291–297.
- (54) Varki, A., and Diaz, S. (1984) The release and purification of sialic acids from glycoconjugates: methods to minimize the loss and migration of O-acetyl groups. *Anal. Biochem.* **137**, 236–247.
- (55) Vanqualef, E., Simon, S., Marquant, G., Garcia, E., Klimerek, G., Delepine, J. C., Cieplak, P., and Dupradeau, F.-Y. (2011) RED Server: a web service for deriving RESP and ESP charges and building force field libraries for new molecules and molecular fragments. *Nucleic Acids Res.* **39**, W511–W517.
- (56) Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W., and Klein, M. L. (1983) Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **79**, 926–935.
- (57) Sattelle, B. M., and Almond, A. (2014) Shaping up for structural glycomics: a predictive protocol for oligosaccharide conformational analysis applied to N-linked glycans. *Carbohydr. Res.* **383**, 34–42.
- (58) Haranczyk, M., and Gutowski, M. (2007) Differences in electrostatic potential around DNA fragments containing guanine and 8-oxo-guanine. *Theor. Chem. Acc.* **117**, 291–296.
- (59) Ren, P. Y., and Ponder, J. W. (2003) Polarizable atomic multipole water model for molecular mechanics simulation. *J. Phys. Chem. B* **107**, 5933–5947.
- (60) Becke, A. D. (1993) Density-functional thermochemistry. 3. The role of exact exchange. *J. Chem. Phys.* **98**, 5648–5652.
- (61) Binkley, J. S., Pople, J. A., and Hehre, W. J. (1980) Self-consistent molecular-orbital methods. 21. Small split-valence basis-sets for 1st-row elements. *J. Am. Chem. Soc.* **102**, 939–947.
- (62) Varki, A., Muchmore, E., and Diaz, S. (1986) A sialic acid-specific O-acetyltransferase in human erythrocytes: possible identity with

esterase D, the genetic marker of retinoblastomas and Wilson disease. *Proc. Natl. Acad. Sci. U. S. A.* 83, 882–886.

(63) Higa, H. H., Diaz, S., and Varki, A. (1987) Biochemical and genetic evidence for distinct membrane-bound and cytosolic sialic acid O-acetyl-esterases: serine-active-site enzymes. *Biochem. Biophys. Res. Commun.* 144, 1099–1108.

(64) Butor, C., Higa, H. H., and Varki, A. (1993) Structural, immunological, and biosynthetic studies of a sialic acid-specific O-acetyl-esterase from rat liver. *J. Biol. Chem.* 268, 10207–10213.

(65) Bakkens, M. J., Zeng, Q., Feitsma, L. J., Hulswit, R. J., Li, Z., Westerbeke, A., van Kuppeveld, F. J., Boons, G. J., Langereis, M. A., Huizinga, E. G., and de Groot, R. J. (2016) Coronavirus receptor switch explained from the stereochemistry of protein-carbohydrate interactions and a single mutation. *Proc. Natl. Acad. Sci. U. S. A.* 113, E3111–3119.

(66) Kamerling, J. P., Schauer, R., Shukla, A. K., Stoll, S., Van Halbeek, H., and Vliegthart, J. F. (1987) Migration of O-acetyl groups in N,O-acetylneuraminic acids. *Eur. J. Biochem.* 162, 601–607.

(67) Diaz, S., Higa, H. H., Hayes, B. K., and Varki, A. (1989) O-Acetylation and de-O-acetylation of sialic acids. 7- and 9-O-acetylation of alpha 2,6-linked sialic acids on endogenous N-linked glycans in rat liver Golgi vesicles. *J. Biol. Chem.* 264, 19416–19426.

(68) Hara, S., Yamaguchi, M., Takemori, Y., Furuhashi, K., Ogura, H., and Nakamura, M. (1989) Determination of mono-O-acetylated N-acetylneuraminic acids in human and rat sera by fluorometric high-performance liquid chromatography. *Anal. Biochem.* 179, 162–166.

(69) Salomon-Ferrer, R., Case, D. A., and Walker, R. C. (2013) An overview of the Amber biomolecular simulation package. *Wiley Interdiscipl. Rev. Comput. Mol. Sci.* 3, 198–210.

(70) Kirschner, K. N., Yongye, A. B., Tschampel, S. M., Gonzalez-Outeirino, J., Daniels, C. R., Foley, B. L., and Woods, R. J. (2008) GLYCAM06: A generalizable biomolecular force field. *Carbohydrates. J. Comput. Chem.* 29, 622–655.

(71) Wang, L.-P., and Song, C. (2016) Geometry optimization made simple with translation and rotation coordinates. *J. Chem. Phys.* 144, 214108.

(72) Ufimtsev, I. S., and Martinez, T. J. (2008) Quantum chemistry on graphical processing units. 1. Strategies for two-electron integral evaluation. *J. Chem. Theory Comput.* 4, 222–231.

(73) Ufimtsev, I. S., and Martinez, T. J. (2009) Quantum chemistry on graphical processing units. 3. Analytical energy gradients, geometry optimization, and first principles molecular dynamics. *J. Chem. Theory Comput.* 5, 2619–2628.

(74) Ufimtsev, I. S., and Martinez, T. J. (2009) Quantum chemistry on graphical processing units. 2. Direct self-consistent-field implementation. *J. Chem. Theory Comput.* 5, 1004–1015.

(75) Wang, J. M., Wolf, R. M., Caldwell, J. W., Kollman, P. A., and Case, D. A. (2004) Development and testing of a general amber force field. *J. Comput. Chem.* 25, 1157–1174.