

# Reversible *O*-Acetyl Migration within the Sialic Acid Side Chain and Its Influence on Protein Recognition

Yang Ji, Aniruddha Sasmal, Wanqing Li, Lisa Oh, Saurabh Srivastava, Audra A. Hargett, Brian R. Wasik, Hai Yu, Sandra Diaz, Biswa Choudhury, Colin R. Parrish, Darón I. Freedberg, Lee-Ping Wang, Ajit Varki,\* and Xi Chen\*



Cite This: *ACS Chem. Biol.* 2021, 16, 1951–1960



Read Online

ACCESS |



Metrics & More




Article Recommendations




Supporting Information

**ABSTRACT:** *O*-Acetylation is a common naturally occurring modification of carbohydrates and is especially widespread in sialic acids, a family of nine-carbon acidic monosaccharides. *O*-Acetyl migration within the exocyclic glycerol-like side chain of mono-*O*-acetylated sialic acid reported previously was from the C7- to C9-hydroxyl group with or without an 8-*O*-acetyl intermediate, which resulted in an equilibrium that favors the formation of the 9-*O*-acetyl sialic acid. Herein, we provide direct experimental evidence demonstrating that *O*-acetyl migration is bidirectional, and the rate of equilibration is influenced predominantly by the pH of the sample. While the *O*-acetyl group on sialic acids and sialoglycans is stable under mildly acidic conditions (pH < 5, the rate of *O*-acetyl migration is extremely low), reversible *O*-acetyl migration is observed readily at neutral pH and becomes more significant when the pH increases to slightly basic. Sialoglycan microarray studies showed that esterase-inactivated porcine torovirus hemagglutinin-esterase bound strongly to sialoglycans containing a more stable 9-*N*-acetylated sialic acid analog, but these compounds were less resistant to periodate oxidation treatment compared to their 9-*O*-acetyl counterparts. Together with prior studies, the results support the possible influence of sialic acid *O*-acetylation and *O*-acetyl migration to host–microbe interactions and potential application of the more stable synthetic *N*-acetyl mimics.



Sialic acids (Sias) are a family of negatively charged nine-carbon monosaccharides typically found at the non-reducing end of glycans on glycoconjugates of vertebrates. Remarkable structural diversity has been observed for Sias, which include *N*-acetylneuraminic acid (Neu5Ac, **1**, Figure 1),



**Figure 1.** Structures of *N*-acetylneuraminic acid (Neu5Ac, **1**) and its 9-*O*-, 8-*O*-, and 7-*O*-acetyl forms (**2–4**).

*N*-glycolylneuraminic acid (Neu5Gc), 2-keto-3-deoxynonulosonic acid (Kdn), and their modified forms.<sup>1–4</sup> *O*-Acetylation of the hydroxyl groups at C9 (**2**), C8 (**3**), C7 (**4**), and/or C4 is the most common Sia modification.

Numerous biological functions of Sias and their involvement in human health and human diseases have been recognized. Many of these roles can be influenced by Sia *O*-acetylation. For example, while *O*-acetylation of the C9 hydroxyl group (C9-OH) in Sia prevents it from being recognized by hemag-

glutinins of influenza A and B viruses, it is required for influenza C and D virus hemagglutinin binding.<sup>5–9</sup> In addition, 7-*O*-acetylation of Sias in group B Streptococcal capsular polysaccharides blocks the binding of Siglec-9 to a greater extent than 9-*O*-acetylation of Sias.<sup>10</sup> The biological implications of *O*-acetylation in Sias have been reviewed and summarized.<sup>1,2,11</sup> Although the importance of Sia *O*-acetylation is increasingly recognized, its functional studies have been hampered by the instability of *O*-acetyl (OAc) groups and their propensity to migrate under biological and analytical conditions. OAc groups are especially labile under basic conditions frequently used in procedures to release, purify, and analyze Sias and sialoglycans.<sup>12</sup> While migration of the Sia 4-*O*-acetyl group was not observed,<sup>13</sup> *O*-acetyl migration within the exocyclic glycerol-like side chain of Sia from C7-OH to C9-OH to reach an equilibrium favoring the formation of 9-OAc-

**Special Issue:** Chemical Glycobiology

**Received:** December 29, 2020

**Accepted:** March 15, 2021

**Published:** March 26, 2021



Sias was reported and was believed to contribute to the prevalence of 9-OAc-Sias among OAc-Sias in nature.<sup>12–15</sup> In sialoglycans where the Sia C9-OH position is already occupied, such as in di-*O*-acetylated Sias or  $\alpha$ 2–9-linked polysialic acids, *O*-acetyl migration from C7-OH to C8-OH in Sia was also observed.<sup>13,14</sup> Studies of sialate:7(9)-*O*-acetyltransferase suggested that Ac was added by the enzyme to the C7-OH of Sia, which then migrated to the C9-OH.<sup>16</sup> An additional Ac could then be added by the enzyme to the C7-OH for the formation of 7,9-di-*O*-acetylated Neu5Ac (Neu5,7,9Ac<sub>3</sub>). Further migration of the Ac group from C7-OH to C8-OH followed by the addition of another Ac group to C7-OH led to the formation of Neu5,7,8,9Ac<sub>4</sub>.<sup>13,16</sup> Alternatively, the hydrolysis of OAc-Sia by sialate-*O*-acetyltransferase-catalyzed reactions was hypothesized to also involve *O*-acetyl migration as a 9-OAc-substrate was the preferred substrate while the OAc group was initially on the C7 of Sia.<sup>17,18</sup>

During recent sialoglycan microarray studies, we found that sialoglycans containing 9-*N*-acetyl-Neu5Ac (Neu5Ac9NAc, a stable synthetic mimic of Neu5,9Ac<sub>2</sub>) showed stronger binding to a Neu5,9Ac<sub>2</sub>-glycan-recognizing protein than their Neu5,9Ac<sub>2</sub>-glycan homologues. The enhancement in its binding to Neu5Ac9NAc-glycans exceeded the gain that would be expected from preventing the de-*O*-acetylation of Neu5,9Ac<sub>2</sub>, indicating that *O*-acetyl migration of Neu5,9Ac<sub>2</sub> in the sialoglycans may contribute to the weaker binding of the protein to the Neu5,9Ac<sub>2</sub>-glycans. To gain additional insight into Sia *O*-acetyl migration and to investigate whether the Ac group at the C9-OH of Sia can undergo reversible migration to the C7-OH and/or C8-OH positions, we improved 1,2-diamino-4,5-methylenedioxybenzene dihydrochloride (DMB) derivatization and high-performance liquid chromatography analysis (DMB-HPLC) by developing a low-temperature pH-controlled derivatization process to minimize *O*-acetyl migration during analysis. Nuclear magnetic resonance (NMR) studies of Neu5,9Ac<sub>2</sub> and Neu5,9Ac<sub>2</sub>-glycan samples at different pH values as well as preparation of deuterium-labeled OAc-Sia and its use for analyzing *O*-acetyl migration at different pHs were carried out. We also performed sialoglycan microarray studies of esterase-inactivated porcine torovirus (PToV) hemagglutinin-esterase (HE) using paired sialoglycan probes containing Neu5,9Ac<sub>2</sub> or its more stable 9-*N*-acetyl analog, Neu5Ac9NAc. The sensitivities of a set of sialoglycans containing Neu5Ac, Neu5,9Ac<sub>2</sub>, or Neu5Ac9NAc to periodate oxidation treatment were also compared. All experimental results were consistent with reversible *O*-acetyl migration at neutral or slightly basic conditions. The potential impact of Sia *O*-acetyl migration on other biomolecular recognition and bioprocess regulation events awaits further exploration.

## RESULTS AND DISCUSSION


**DMB-HPLC Sialic Acid Quantitative Assay Improvement to Minimize *O*-Acetyl Migration during Analysis.** DMB derivatization followed by HPLC analysis (DMB-HPLC) has been commonly used to quantify Sias.<sup>12</sup> Conventional DMB derivatization (50 °C for 2.5 h) was successful for non-*O*-acetylated Sias but led to a significant loss of OAc groups and unwanted *O*-acetyl migration in OAc-Sias. Lowering the reaction temperature to 4 °C significantly decreased the level of de-*O*-acetylation or *O*-acetyl migration, but concomitantly decreased DMB derivatization efficiencies even with extended reaction times (e.g., 48 h) for samples originally prepared in a buffer at neutral or basic pH. To

quantify OAc-Sias more accurately with minimal OAc loss or migration while retaining optimal DMB derivatization yields, an improved DMB derivatization method was developed. Controlling the pH during sample preparation for DMB derivatization at 4 °C was critical to achieving optimal yields. Previous low-temperature derivatization procedures used DMB in 0.02 M trifluoroacetic acid.<sup>19</sup> DMB reagent, mixed with the same volume of a sample prepared in water (pH 7.0) or sodium acetate solution (50 mM, pH = 5.0 or 7.0), resulted in samples with varied pH values of 2.7, 4.3, and 4.7, respectively. The pH variation may have caused the inconsistencies in derivatization efficiencies. To maintain a consistent acidic pH range for optimal DMB derivatization of  $\alpha$ -keto acids such as Sias, the DMB reagent was prepared in a high concentration (1.6 M) acetic acid solution. Mixing with the same volume of Sia solution in either water or in a 50 mM sodium acetate solution of pH = 5.0 or 7.0, yielded samples with a steady pH range (3.8–4.1). Improved peak separation in HPLC-based analysis of DMB-derivatized Sias was further optimized by changing the isocratic conditions (acetonitrile/methanol/water = 8:7:85, v/v) to a linear gradient (acetonitrile/methanol/water from 7:7:86 to 11:7:82, v/v) in a 75 min run.

Figure 2A shows that DMB derivatization at 4 °C for 48 h with improved sample preparation conditions better preserved the OAc groups on Sias and reduced de-*O*-acetylation and *O*-acetyl migration compared to the reaction performed at 50 °C for 2 h. Additionally, consistent derivatization efficiencies and results were obtained when the DMB reagent was prepared in acetic acid (1.6 M) and mixed with Sia in the same volume in a cold sodium acetate solution (50 mM) of a different pH (pH 5.0 or pH 7.0) at 4 °C for 48 h (Figure 2B). A Neu5,9Ac<sub>2</sub> sample synthesized by regioselective acetylation of Neu5Ac using trimethyl orthoacetate with purity confirmed by NMR analysis previously<sup>21</sup> was converted to a mixture containing Neu5Ac (~3%), Neu5,7Ac<sub>2</sub> (~2%), Neu5,8Ac<sub>2</sub> (~30%), and Neu5,9Ac<sub>2</sub> (~65%) after roughly three years of storage as a lyophilized powder at –20 °C.

**pH-Dependent Reversible *O*-Acetyl Migration among the Hydroxyl Groups on the Glycerol Side Chain of Neu5Ac by DMB-HPLC and NMR Studies.** The pH-dependent *O*-acetyl migration in OAc-Sias was initially observed for samples prepared from bovine submaxillary mucins by gas liquid chromatograph (GLC) analysis<sup>12</sup> and was later confirmed by more thorough HPLC and <sup>1</sup>H NMR spectroscopy studies.<sup>13</sup> As the samples used for the studies had high percentages of 7/8-OAc-Sias, the *O*-acetyl migration observed was from C7-OH or C8-OH to C9-OH. Therefore, *O*-acetyl migration in mono-OAc-Sia was assumed to be unidirectional with a rate dependent on both pH and temperature. Upon reaching equilibrium after an extended incubation period (>1250 min, 37 °C, pH 7.2), about 10% Neu5,7Ac<sub>2</sub> remained by <sup>1</sup>H NMR analysis.<sup>13</sup> Neu5,8Ac<sub>2</sub> was not detected, therefore whether it was involved in the migration was unresolved in previous studies.


Using the synthetic Neu5,9Ac<sub>2</sub> sample that was converted to a mixture of Neu5Ac (~3%), Neu5,7Ac<sub>2</sub> (~2%), Neu5,8Ac<sub>2</sub> (~30%), and Neu5,9Ac<sub>2</sub> (~65%) after approximately three years of storage as a lyophilized powder at –20 °C, we revisited Sia *O*-acetyl migration using the improved DMB-HPLC method described above. When the sample (starting material 1 or SM1; Figure 3) was incubated in a sodium acetate solution with a pH in the range of 3.0–8.0 at 37 °C for 2 h, *O*-acetyl migration from C8-OH to C9-OH was clearly



**Figure 2.** Comparison of DMB derivatization-HPLC analysis methods for quantification of OAc-Sias. (A) HPLC chromatograms of a DMB-derivatized water-dissolved OAc-Sia sample at 4 °C for 48 h (top panel) and at 50 °C for 2 h (bottom panel). (B) HPLC chromatograms of a DMB-derivatized OAc-Sia sample (at 4 °C for 48 h) by mixing the DMB reagent in acetic acid (1.6 M) with the same volume of the OAc-Sia sample in a cold sodium acetate solution (50 mM) with a pH of 5.0 (top panel) or 7.0 (bottom panel). Sias from bovine submaxillary mucin (BSM) previously studied by liquid chromatography-electrospray ionization-mass spectrometry (LC-ESI-MS)<sup>20</sup> were used as the standards.

observed at pH  $\geq$  5.0, although the degree of migration was larger at pH  $\geq$  7.0. For example, at pH 8.0 where de-*O*-acetylation was less than 1%, Neu5,8Ac<sub>2</sub> decreased from 32.4% to 11.8% while Neu5,9Ac<sub>2</sub> concomitantly increased from 65.6% to 85.1%. A slight increase (2.0% to 3.0%) of Neu5,7Ac<sub>2</sub> was also observed.

In another more freshly purified Neu5,9Ac<sub>2</sub> sample (starting material 2 or SM2), incubation in a sodium acetate solution at pH 8.0 and 37 °C for 2 h increased the percentages of Neu5,7Ac<sub>2</sub> and Neu5,8Ac<sub>2</sub> from 0.8% to 3.1% and 6.1% to 10.7%, respectively, while the percentage of Neu5,9Ac<sub>2</sub> decreased from 93.1% to 86.2% (Figure 3). The results demonstrated *O*-acetyl migration from C9-OH to C8-OH and C7-OH, which is the opposite direction of that reported previously for mono-*O*-acetylated Sias. Similar to SM1, *O*-acetyl migration in SM2, although in the opposite direction to that observed for SM1, occurred at pH  $\geq$  5.0 and at a higher level at pH  $\geq$  7.0. The migration in both directions seen in SM1 and SM2 seemed to have reached an equilibrium among



**Figure 3.** *O*-Acetyl migration in OAc-Sia monosaccharide samples. DMB-HPLC analysis of *O*-acetyl migration with three OAc-Sia monosaccharide samples, each containing different amounts of Neu5,7Ac<sub>2</sub> (black columns), Neu5,8Ac<sub>2</sub> (white columns), and Neu5,9Ac<sub>2</sub> (gray columns). OAc-Sia samples were incubated at different pHs at 37 °C for 2 h. Before incubation, starting material 1 (SM1) contains 2.0% of Neu5,7Ac<sub>2</sub>, 32.4% of Neu5,8Ac<sub>2</sub>, and 65.6% of Neu5,9Ac<sub>2</sub>; starting material 2 (SM2) contains 0.8% of Neu5,7Ac<sub>2</sub>, 6.1% of Neu5,8Ac<sub>2</sub>, and 93.1% of Neu5,9Ac<sub>2</sub>; starting material 3 (SM3) contains 1.4% of Neu5,7Ac<sub>2</sub>, 8.5% of Neu5,8Ac<sub>2</sub>, and 90.1% of Neu5,9Ac<sub>2</sub>.

Neu5,9Ac<sub>2</sub> (~87%), Neu5,8Ac<sub>2</sub> (~10%), and Neu5,7Ac<sub>2</sub> (3%) at pH 8.0. When the sample pH was lowered to 3.0, no significant *O*-acetyl migration was observed for over 2 h. Similar results were obtained for an OAc-Sia sample (starting material 3 or SM3), which contained 90.1% Neu5,9Ac<sub>2</sub>, 8.5% Neu5,8Ac<sub>2</sub>, and 1.4% Neu5,7Ac<sub>2</sub> before incubation, and a mixture of 88% Neu5,9Ac<sub>2</sub>, 10% Neu5,8Ac<sub>2</sub>, and 2% Neu5,7Ac<sub>2</sub> was obtained after incubation at pH 7.0 and 37 °C for 2 h.


To determine the experimental errors of the DMB-HPLC assays (e.g., Figure 3), the assays were repeated for both SM1 and SM3 in triplicate. The standard deviations of the percentages of individual OAc-Sia determined did not exceed 0.2% (Table S1).

SM3 was also used in a series of two-dimensional NMR experiments including <sup>1</sup>H–<sup>13</sup>C HSQC,<sup>22</sup> HSQC-TOCSY,<sup>23,24</sup> and long-range (LR)-HSQMBC<sup>25</sup> (Figures S1–S2; 37 °C, D<sub>2</sub>O, pH\* 6.14) to assign the chemical shifts of *O*-acetylated Sias, including the  $\alpha$  and  $\beta$  anomers of Neu5,9Ac<sub>2</sub>, Neu5,8Ac<sub>2</sub>, Neu5,7Ac<sub>2</sub>, and Neu5Ac (Table S2). The LR-HSQMBC showed a cross peak at <sup>1</sup>H 4.9 ppm and <sup>13</sup>C 173.3 ppm, confirming *O*-acetyl at C8 in Neu5,8Ac<sub>2</sub> (Figure S2). Previously, a similar <sup>1</sup>H peak at  $\delta$  = 5.1 ppm which displayed a doublet of doublets of doublets splitting pattern (ddd) was assigned to the H8 of Neu5,8,9Ac<sub>3</sub> at pH\* 7.5.<sup>13</sup> The assignments were then used to interpret one-dimensional <sup>1</sup>H NMR spectra of another sample of SM3 recorded under acidic (pH\* 3.5) conditions at  $t$  = 0 min and after incubation at 37 °C for 120 min and 12 h. The H9s of Neu5,9Ac<sub>2</sub> were integrated as 1.00 for reference. The ddd peak at  $\delta$  = 4.8 ppm for the H-8 of Neu5,8Ac<sub>2</sub> and the dd peak at  $\delta$  = 4.9 for the H-7 of Neu5,7Ac<sub>2</sub> were used for integration to determine the percentages of individual species. Approximately 10% of Neu5,8Ac<sub>2</sub> ( $\delta$  = 4.95 ppm) and 2.1% of Neu5,7Ac<sub>2</sub> ( $\delta$  = 5.05 ppm) were detected in the sample at pH\* 3.5 at the beginning ( $t$  = 0 min), which remained unchanged throughout the NMR experiment (Figure S3A–C). The pH\* of the sample was then adjusted to 7.0, and the spectra were recorded again at  $t$  = 0 min and after incubation at 37 °C for 120 min and 12 h. After the pH\* was changed to 7.0, about 12% of

Neu5,8Ac<sub>2</sub> and 3% of Neu5,7Ac<sub>2</sub> were detected at  $t = 0$  min, which remained unchanged during the incubation at 37 °C for 12 h (Figure S3D–F). The results indicated that the sample prepared in an aqueous solution at pH\* 7.0 reached an equilibrium for *O*-acetyl migration at the beginning of the NMR experiment and the ratios of OAc-Sias agreed with those obtained by the improved DMB-HPLC method described above.

**Confirmation of Simultaneous Bidirectional *O*-Acetyl Migration in OAc-Sias by Synthesizing a Deuterated Neu5,9Ac<sub>2</sub>.** To confirm the reversible *O*-acetyl migration in OAc-Sias, Neu5,9Ac<sub>2</sub> with a 9-*O*-deuterated acetyl group (deuterated Neu5,9Ac<sub>2</sub>) was synthesized by modifying a lipase-catalyzed *O*-acetylation reaction using deuterated vinyl acetate as the acetylation reagent.<sup>26</sup> The deuterated Neu5,9Ac<sub>2</sub> sample was mixed with nondeuterated Neu5,8Ac<sub>2</sub> and Neu5,9Ac<sub>2</sub>, and an OAc-Sia mixture containing 87.3% Neu5,9Ac<sub>2</sub> (64.0% nondeuterated and 23.3% deuterated), 12.2% Neu5,8Ac<sub>2</sub> (9.2% nondeuterated and 3.0% deuterated), and 0.5% Neu5,7Ac<sub>2</sub> (nondeuterated) was obtained, as confirmed by LC-MS (the control sample in Table S3 and Figure S4). The effective separation of Neu5,7Ac<sub>2</sub>, Neu5,8Ac<sub>2</sub>, and Neu5,9Ac<sub>2</sub> by HPLC and comparison with bovine submaxillary mucin (BSM) Sia standards ensured the correct designation of OAc-Sia species. The differentiation and quantification of the deuterated and nondeuterated forms of Neu5,9Ac<sub>2</sub>, Neu5,8Ac<sub>2</sub>, and Neu5,7Ac<sub>2</sub> were based on the peaks in the extracted ion current (XIC) chromatograms of the MS. If Sia *O*-acetyl migration were unidirectional from C7 and/or C8 to C9-OH, we would see an unchanged percentage of each OAc-Sia species, or a slight decrease of Neu5,8Ac<sub>2</sub> (both deuterated and nondeuterated) in the given OAc-Sia mixture. However, in a dynamic equilibrium, the deuterated Ac group on deuterated Neu5,9Ac<sub>2</sub> would migrate back to form [8-deuteroacetyl]-Neu5,8Ac<sub>2</sub>, while the nondeuterated acetyl group on Neu5,8Ac<sub>2</sub> would migrate to C9-OH concurrently. Therefore, we would observe an increase in both [8-deuteroacetyl]-Neu5,8Ac<sub>2</sub> and nondeuterated Neu5,9Ac<sub>2</sub>.


As shown in Figure 4, the net percent increase in deuterated Neu5,8Ac<sub>2</sub> by incubation at 7.0 or 8.0 (from 3.0% to 4.6% and 5.1%, respectively) clearly demonstrated the migration of the deuterated Ac group from C9-OH to C8-OH (Table S3). The



**Figure 4.** Concurrent interconversion between Neu5,8Ac<sub>2</sub> and Neu5,9Ac<sub>2</sub> at neutral or slightly basic pH. A [9-deuteroacetyl]-Neu5,9Ac<sub>2</sub> sample was mixed with nondeuterated Neu5,8Ac<sub>2</sub>/Neu5,9Ac<sub>2</sub> and incubated at various pHs at 37 °C for 30 min followed by LC-MS analysis. The starting mixture without incubation was used as a control (ctrl). The percentage of deuterated and nondeuterated Neu5,8Ac<sub>2</sub> (A) and Neu5,9Ac<sub>2</sub> (B) was calculated based on the corresponding peak areas in the XIC chromatogram. Deuterated samples are shown in black columns, and nondeuterated samples are shown in gray columns.

net percent decrease in nondeuterated Neu5,8Ac<sub>2</sub> under the same conditions (from 9.2% to 5.1% at pH 7.0, to 4.4% at pH 8.0) indicated the migration of nondeuterated Neu5,8Ac<sub>2</sub> from C8-OH to C9-OH. Furthermore, the fact that 0.3% deuterated Neu5,7Ac<sub>2</sub>, which was absent in the control, was detected at pH 8.0 together with the consistent results in Figure 3, suggests the involvement of the C7-OH position in the bidirectional *O*-acetyl migration.

To further investigate the participation of C7-OH in OAc-Sia *O*-acetyl migration, the deuterated Neu5,9Ac<sub>2</sub> sample was mixed with nondeuterated Neu5,7Ac<sub>2</sub> and Neu5,9Ac<sub>2</sub>, and a mixture of 69.9% Neu5,9Ac<sub>2</sub> (56.0% deuterated and 13.9% nondeuterated), 3.7% Neu5,8Ac<sub>2</sub> (2.3% deuterated and 1.4% nondeuterated), and 26.4% Neu5,7Ac<sub>2</sub> (all nondeuterated) was obtained (the control sample in Table S4). *O*-Acetyl migration was significant at pH 7.0 and 8.0 (Figure 5 and




**Figure 5.** Concurrent interconversion between Neu5,7Ac<sub>2</sub> and Neu5,9Ac<sub>2</sub> at neutral or slightly basic pH. A [9-deuteroacetyl]-Neu5,9Ac<sub>2</sub> was mixed with nondeuterated Neu5,7Ac<sub>2</sub>/Neu5,9Ac<sub>2</sub> and incubated at various pHs at 37 °C for 2 h followed by LC-MS analysis. The starting mixture without incubation was used as a control. The percentages of deuterated and nondeuterated Neu5,7Ac<sub>2</sub> (A), Neu5,8Ac<sub>2</sub> (B), and Neu5,9Ac<sub>2</sub> (C) were calculated based on the corresponding peak areas in the XIC chromatogram. Deuterated samples are shown in black columns, and nondeuterated samples are shown in gray columns.

Table S4). The net percent increases in deuterated Neu5,7Ac<sub>2</sub> and deuterated Neu5,8Ac<sub>2</sub> by incubation at pH 7.0 or 8.0 demonstrated migration of the deuterated Ac from C9-OH to C7-OH and C8-OH. The net percent decrease of nondeuterated Neu5,7Ac<sub>2</sub> and the net percent increases of nondeuterated Neu5,8Ac<sub>2</sub> and Neu5,9Ac<sub>2</sub> clearly demonstrated the migration of Ac from C7-OH to C8-OH and C9-OH.

To determine the experimental errors in the LC-MS assays (e.g., Figures 4 and 5), the study was repeated for SM3 in triplicate. The standard deviations of the percentages of individual OAc-Sia determined ranged between 0.02–0.09% (Table S5).

**9-OAc in Neu5,9Ac<sub>2</sub>-Containing Sialoglycans Can Also Migrate to C8-OH and C7-OH in a pH-Dependent Manner.** Using two structurally defined synthetic sialoglycans Neu5,9Ac<sub>2</sub>α3Galβ3GlcNAcβProNH<sub>2</sub> and Neu5,9Ac<sub>2</sub>α3Galβ4GlcNAcβProNH<sub>2</sub>, pH-dependent *O*-acetyl

migration in glycosidically bound OAc-Sias was also analyzed. After the sialoglycans were incubated at various pHs in a water bath at 37 °C for 2 h, the terminal Sias were quickly released by incubating each sample with 10 mU of *Arthrobacter ureafaciens* sialidase for 10 min before low temperature DMB-HPLC analysis of OAc-Sia composition. The purpose of the relatively short incubation time with the sialidase was to minimize *O*-acetyl migration during sample analysis. The increase in pH accelerated migration of the Ac group from C9-OH to C8-OH in Sia (Figure 6). Neu5,8Ac<sub>2</sub> increased from



**Figure 6.** *O*-Acetyl migration on OAc-Sia-containing sialoglycans. Neu5,9Ac<sub>2</sub>α3Galβ3GlcNAcβProNH<sub>2</sub> (A) and Neu5,9Ac<sub>2</sub>α3Galβ4GlcNAcβProNH<sub>2</sub> (B) at a different pH by comparing the composition of Neu5,9Ac<sub>2</sub> (gray columns), Neu5,8Ac<sub>2</sub> (white columns), and Neu5,7Ac<sub>2</sub> (black columns) released by sialidase treatment before the low temperature DMB-HPLC analysis.

1.7% at pH 5.0 to 8.0% at pH 8.0 in Neu5,9Ac<sub>2</sub>α3Galβ3GlcNAcβProNH<sub>2</sub>. For Neu5,9Ac<sub>2</sub>α3Galβ4GlcNAcβProNH<sub>2</sub>, Neu5,8Ac<sub>2</sub> increased from 0.4% at pH 5.0 to 7.0% at pH 8.0. A similar increase in %Neu5,8Ac<sub>2</sub> was observed for Sias on bovine submaxillary mucin (from 0.2% at pH 5.0 to 4.9% at pH 8.0) although the results were complicated by the accompanying loss of di-OAc-Sias (Figure S5).

The pH-dependent *O*-acetyl migration from C9-OH to C8 and C7-OH in sialoglycans, such as Neu5,9Ac<sub>2</sub>α3Galβ4GlcβProN<sub>3</sub>, was also confirmed by NMR studies at pH\* 3.5 and pH\* 8.0. In this study, one of the two diastereotopic protons on C9 of Neu5,9Ac<sub>2</sub> (dd, δ 4.25 ppm), the proton on C8 of Neu5,8Ac<sub>2</sub> (ddd, δ 5.2 ppm), and the proton on C7 of Neu5,7Ac<sub>2</sub> (dd, δ 5.1 ppm)<sup>27,28</sup> were used for peak integration, the total area under which was normalized to 1.00 (Figure S6). The integrals of the H3-equatorial and H3 axial signals were also used as references as the *O*-acetyl substitution on C9-OH, C8-OH, or C7-OH of Neu5Ac in sialoglycans only had minimal influence on the chemical shift of H3 in Neu5Ac.<sup>29</sup> Incubating the sample in D<sub>2</sub>O at pH\* 3.5 for 2 h (entries 1 and 2) did not change the ratios of OAc-Sias in the sialoglycans (~92% Neu5,9Ac<sub>2</sub>-glycan, 5.8–5.9% Neu5,8Ac<sub>2</sub>-glycan, and 2.5% Neu5,7Ac<sub>2</sub>-glycan, Table 1). The presence of Neu5,8Ac<sub>2</sub>-glycan and Neu5,7Ac<sub>2</sub>-glycan in the initial time point (entry 1) at *t* = 0 may be due to *O*-acetyl migration that occurred during sample preparation. The ratio of different OAc-Sias in the sialoglycans changed quickly to 83% Neu5,9Ac<sub>2</sub>-glycan, 7.6% Neu5,8Ac<sub>2</sub>-glycan, and 9.2% Neu5,7Ac<sub>2</sub>-glycan when the pH\* of the sample was adjusted to 8.0 (entry 3 in Table 1, Figure S6C). The ratio remained constant during a 2 h incubation at pH\* 8.0 and 37 °C (entry 4 in Table 1, Figure S6D), indicating that an equilibrium was established shortly after the pH\* was adjusted to 8.0, before the first NMR spectrum was taken at *t* = 0 min. Changing the pH\* of the sample back to 3.5 (entry 5 in Table 1, Figure S6E)

**Table 1.** pH\*-Dependent *O*-Acetyl Migration in Neu5,9Ac<sub>2</sub>α3Galβ4GlcβProN<sub>3</sub> Observed by Percent Changes in Individual Mono-*O*-acetyl Sia Species by NMR Studies (the Corresponding NMR Spectra Are in Figure S6)

no.	pH* (time)	Neu5,9Ac <sub>2</sub>	Neu5,8Ac <sub>2</sub>	Neu5,7Ac <sub>2</sub>	total
1	3.5 ( <i>t</i> = 0)	0.9160	0.0589	0.0250	1.00
2	3.5 ( <i>t</i> = 2 h)	0.9175	0.0579	0.0246	1.00
3	8.0 ( <i>t</i> = 0)	0.8325	0.0758	0.0917	1.00
4	8.0 ( <i>t</i> = 2 h)	0.8335	0.0758	0.0907	1.00
5	3.5 ( <i>t</i> = 0)	0.8322	0.0760	0.0919	1.00
6	3.5 ( <i>t</i> = 3 d)	0.8339	0.0746	0.0915	1.00
7	8.0 ( <i>t</i> = 0)	0.8274	0.0766	0.0925	1.00
8	8.0 ( <i>t</i> = 12 h)	0.8314	0.0763	0.0923	1.00


did not alter the ratio of OAc-Sias in the sialoglycans significantly, even after a 3-day incubation at 37 °C (entry 6 in Table 1, Figure S6F). This is consistent with the observations for unbound OAc-Sia monosaccharides described above. Adjusting the pH\* of the sample again to pH\* 8.0 (entry 7 in Table 1, Figure S6G) did not significantly alter the ratio of glycosidically bound OAc-Sias as the equilibrium was reached previously. The ratio remained unchanged after an additional 12 h at 37 °C (entry 8 in Table 1, Figure S6H).

#### A Neu5,9Ac<sub>2</sub>-Binding Protein, PToV HE, Shows Stronger Binding to Immobilized Sialoglycans Containing Neu5Ac9NAc than Those Containing Neu5,9Ac<sub>2</sub>.

Our previous glycan microarray study with a matching set of α2–3-linked sialyl lactosides (Siaα3Galβ4GlcβProNH<sub>2</sub>) containing a terminal Neu5Ac, Neu5,9Ac<sub>2</sub>, or its more stable *N*-acetyl analog Neu5Ac9NAc showed that PToV HE recognizes both Neu5,9Ac<sub>2</sub>-glycan and its Neu5Ac9NAc-analog but not Neu5Ac-glycan.<sup>30</sup> Using a library of α2–3- and α2–6-linked synthetic sialoglycans containing different internal glycans and the matching terminal Neu5,9Ac<sub>2</sub> and its Neu5Ac9NAc analog,<sup>31</sup> sialoglycan microarray studies showed that the esterase-inactivated PToV HE bound to Neu5Ac9NAc-glycans with much stronger signals than Neu5,9Ac<sub>2</sub>-glycans (Figure 7). Strong binding signals were also shown for other Neu5Ac9NAc-glycans with unmatched internal glycan structures (the last six bars shown in Figure 7). The lack of binding to Neu5Ac-glycans observed previously was also confirmed, consistent with the report of PToV HE as a Neu5,9Ac<sub>2</sub>-recognizing probe.<sup>32</sup> The lower binding to Neu5,9Ac<sub>2</sub>-glycans, partly due to the lower stability of the 9-OAc group in Neu5,9Ac<sub>2</sub>-glycans compared to the 9-*N*-acetyl group in Neu5Ac9NAc-glycans, may be explained by the *O*-acetyl migration in Neu5,9Ac<sub>2</sub>-glycans. Our study also indicates that the terminal Sia is the main recognition component of PToV HE while the sialyl linkage and the internal glycan structures do not contribute significantly to binding. The study also supports the role of Neu5Ac9NAc-glycans as useful stable mimics of Neu5,9Ac<sub>2</sub>-glycans and potent probes for research.


#### Neu5Ac9NAc-Glycans Are More Sensitive to Periodate Oxidative Cleavage than Neu5,9Ac<sub>2</sub>-Glycans.

Periodate oxidative cleavage of C–C bonds with vicinal hydroxyl groups such as those in carbohydrates is well-known.<sup>33</sup> Selective cleavage of the C–C bonds in the glycerol chain of Sia in sialoglycans to generate an aldehyde group at C7 without affecting other sugars can be achieved by using low concentrations of periodate. The strategy has been successfully used for selective labeling of the sialoglycoproteins on living cell surfaces.<sup>34–37</sup>



**Figure 7.** Sialoglycan microarray binding studies of PToV HE toward Neu5Ac, Neu5,9Ac<sub>2</sub>, and Neu5Ac9NAc-containing glycans. Neu5Ac (black columns), Neu5,9Ac<sub>2</sub> (white columns), and Neu5Ac9NAc-glycans (gray columns) with matched underlying structures were compared. Neu5Ac9NAc-glycans that did not have comparable Neu5Ac or Neu5,9Ac<sub>2</sub>-glycans with matching underlying structures were grouped together and plotted on the right side of the chart. Biotinylated-PToV HE (30 μg mL<sup>-1</sup>) was applied to all glycans.

We found that mild periodate treatment could differentiate the matching set of Siaα3Galβ3(Fucα4)GlcNAcβProNH<sub>2</sub> containing a terminal Neu5Ac, Neu5,9Ac<sub>2</sub>, or its more stable *N*-acetyl analog Neu5Ac9NAc.<sup>38</sup> Among these, the Neu5Ac-glycan was the most sensitive toward periodate treatment. Incubation of samples with 6 mM NaIO<sub>4</sub> at 20 °C for 30–120 min followed by reduction with 30 mM NaBH<sub>4</sub> at 20 °C for 20 min, 2 M AcOH release of the Sia at 80 °C for 1 h, DMB derivatization, and HPLC analysis showed that the Neu5Ac in the Neu5Ac-glycan ( $m/z = 876.34$ ; Figure S8A) was completely converted to its C7-aldehyde derivative by periodate oxidative cleavage in 30 min. Consistent with a previous report,<sup>39</sup> the DMB derivatized C7-Neu5Ac derivative was eluted right after Neu5Ac as shown in the HPLC chromatogram (Figure 8A). The formation of the C7-aldehyde derivative of the sialoglycan after periodate oxidative cleavage was also validated by LC-MS analysis of its borohydride reduced product ( $m/z = 818.34$ ; Figure S8B). Interestingly, Neu5Ac9NAc-glycan (Figure 8B) was less resistant to the periodate oxidative cleavage at C7 than its Neu5,9Ac<sub>2</sub>-glycan counterpart (Figure 8C). While 74.2% of the Neu5,9Ac<sub>2</sub>-glycan was retained after 2 h of periodate treatment (Figure S9B) of a sample originally containing 84.6% Neu5,9Ac<sub>2</sub>-glycan and 15.4% de-*O*-acetylated Neu5Ac-glycan (the *O*Ac loss occurred during storage), only 62.8% of the Neu5Ac9NAc-glycan was retained from a pure starting material under the same conditions (Figure S9A). The C7 derivative ( $m/z = 818.34$ ; Figure S10B) formed from the periodate-treated Neu5Ac9NAc-glycan followed by reduction ( $m/z = 917.37$ ; Figure S10A) was the same as that from the periodate-treated Neu5Ac-glycan followed by reduction ( $m/z = 818.34$ ; Figure S8B). Another experiment using a matching set of Siaα3Galβ4GlcβProNH<sub>2</sub> under the same periodate treatment and downstream assay conditions also showed higher sensitivity of Neu5Ac9NAc-glycan than Neu5,9Ac<sub>2</sub>-glycan. As shown in Figure S11 by DMB-HPLC analysis, while 26.2% Neu5,9Ac<sub>2</sub>-glycan remained after 2 h of periodate treatment of



**Figure 8.** DMB-HPLC assay results of sialoglycan Siaα3Galβ3-(Fucα4)GlcNAcβProNH<sub>2</sub> containing Neu5Ac (A), Neu5Ac9NAc (B), or Neu5,9Ac<sub>2</sub> (C) treated with 6 mM NaIO<sub>4</sub> at pH 6.5 and 20 °C for 30–120 min, followed by sodium borohydride (30 mM) reduction at 20 °C for 20 min and acetic acid (2 M) releasing of Sias at 80 °C for 1 h.

an original sample containing 39.4% Neu5,9Ac<sub>2</sub>-glycan, 62.8% of Neu5Ac9NAc-glycan remained from a pure sample to the same treatment. The data strongly suggest that *O*-acetyl migration may play a role in the resistance of Neu5,9Ac<sub>2</sub> to periodate oxidative cleavage. It was shown previously that amino group substitution of one of the vicinal diols activated the periodate oxidative cleavage of the corresponding C–C bond and *N*-acetylation greatly reduced the reactivity.<sup>40</sup> Nevertheless, the difference of C9-Nac and C9-OAc on the

influence of periodate oxidative cleavage of the diol-containing C7–C8 bond is unclear.

## CONCLUSIONS

In summary, we developed an improved DMB-HPLC method with DMB derivatization carried out at a consistent acidic pH range (3.8–4.1) and a low temperature (4 °C) that allowed quantification of Neu5Ac and its mono-*O*-acetylated derivatives with minimal *O*-acetyl migration or de-*O*-acetylation. Advanced NMR methods (i.e., LR-HSQMBC, etc.)<sup>24</sup> and equipment (cryoprobe, higher fields) used here enabled us to obtain data for lower abundance species that were not observed previously, such as Neu5,8Ac<sub>2</sub>.<sup>13</sup> In addition to the previous observation of *O*-acetyl migration in Sia from C7-OH to C9-OH, we provide direct evidence here that *O*-acetyl group migration in free and glycosidically bound Sia occurs in all directions among C7-OH, C8-OH, and C9-OH in a pH-dependent manner. While *O*-acetyl migration was minimal at pH lower than 5.0, it increased with increasing pH. The synthetic monosaccharide and/or sialoglycan samples with or without deuterium-labeling containing varied percentages of Neu5,9Ac<sub>2</sub>, Neu5,8Ac<sub>2</sub>, and Neu5,7Ac<sub>2</sub> were critical for the observation of bidirectional *O*-acetyl migration. *O*-Acetyl migration may play a role in the higher resistance of Neu5,9Ac<sub>2</sub>-glycans to periodate oxidative cleavage compared to their Neu5Ac9NAc-glycan counterpart. It may also influence the interaction of Neu5,9Ac<sub>2</sub>-glycans with sialoglycan-binding proteins such as a Neu5,9Ac<sub>2</sub>-recognizing esterase-inactivated porcine hemagglutinin-esterase shown here which had a stronger binding to Neu5Ac9NAc-glycans than their Neu5,9Ac<sub>2</sub>-glycan counterparts. Further investigations are needed to fully understand the biological implications of bidirectional *O*-acetyl migration in Sia and sialoglycans.

## METHODS

**O-Acetylated Sia Monosaccharide Preparation.** Trimethyl orthoacetate (0.165 mL, 1.30 mmol) and *p*-TsOH·H<sub>2</sub>O (10 mg) were added to a solution of Neu5Ac (0.2 g, 0.65 mmol) in DMSO (2 mL). The reaction mixture was stirred for 1 h at RT, and then applied to an anion exchange resin column chromatograph (Dowex 1-X8, HCO<sub>2</sub><sup>-</sup> form). The column was washed with water, and the product was eluted with 1 M formic acid. The fractions containing Neu5,9Ac<sub>2</sub> were combined and concentrated. The resulting material was loaded on a Biogel P-2 fine column to obtain the pure product. This compound was stored as the acid form for years, which produced a significant amount of Neu5,8Ac<sub>2</sub> (starting material 1 or SM1). Starting material 2 (SM2) was obtained by releasing from a Neu5,9Ac<sub>2</sub>-containing glycan with 10 mU of *Arthrobacter ureafaciens* sialidase (AUS) in 50 mM sodium acetate (pH 5.5). To obtain starting material 3 (SM3), SM1 was dissolved in H<sub>2</sub>O (1 mL) and purified by HPLC using a reverse-phase C18 column (Phenomenex, 21.2 × 250 mm) with a flow rate of 10 mL min<sup>-1</sup> and a gradient elution of 0–100% acetonitrile in water over 20 min. Mobile phase A, water; mobile phase B, acetonitrile; gradient, 0% B for 3 min, 0% to 100% B over 12 min, 100% B for 2 min, then 100% to 0% B over 3 min. HPLC eluents were monitored by absorption at 210 nm, and glycan-containing fractions were analyzed by TLC and MS. The fractions containing the pure product were collected and lyophilized to obtain pure Neu5,9Ac<sub>2</sub>. Neu5,7Ac<sub>2</sub> was obtained by releasing from the hyper-*O*-acetylated capsular polysaccharides (CPS) of the group B streptococcus COH1 ΔNeuA + N301A NeuA strain. The culture and releasing conditions were the same as described previously.<sup>18</sup> For lipase-catalyzed synthesis of [9-deuteroacetyl]Neu5,9Ac<sub>2</sub>, deuterated vinyl acetate (0.1 mL, 99.2%-D<sub>6</sub>, CDN Isotopes, Lot. DE169) was added with 50 μL of Neu5Ac in pyridine (50 mM), 0.35 mL acetonitrile and 1 mg lipase (from *Pseudomonas* sp., Type XIII, 26

units mg<sup>-1</sup>). The reaction was carried out at 45 °C for 3 days. The organic solvent was removed by nitrogen blow down. The product was then dissolved in ice-cold water and passed through 10 kDa Amicon filter to remove the lipase. The flow-through was lyophilized. Deuterated Neu5,9Ac<sub>2</sub> product was confirmed and quantified by DMB-HPLC and LC-MS as described above (Figure S7).

**Sialoglycans Microarray Study.** The production of the Neu5,9Ac<sub>2</sub>-recognizing probe by use of ectodomains of the Hemagglutinin-Esterase (HE) of porcine torovirus was described in previous work.<sup>41</sup> In this work, an Avi-tag was inserted between the Fc domain and 6X-His and biotinylated according to the suggestions by manufacturers (NEB, Avidity). Biotinylated proteins were concentrated and the buffer was exchanged to PBS. The sialoglycan microarray experiment was performed following the earlier reported literature with a slight modification.<sup>42,43</sup> Chemoenzymatically synthesized sialoglycans were quantitated utilizing an improved low temperature DMB-HPLC method described before. Then, a 10 mM stock solution was prepared (in water) and further diluted to 100 μM in 300 mM Na-phosphate buffer (pH 8.4). Diluted sialoglycans were printed in quadruplets on NHS-functionalized glass slides (PolyAn 3D-NHS; catalog# PO-10400401) using an ArrayIt SpotBot Extreme instrument, dried in the printing chamber, and then blocked using 0.05 M ethanolamine solution in 0.1 M Tris-HCl (pH 9.0). After washing with warm Milli-Q water and centrifuge drying, the slides were stored at 4 °C, under dark in a vacuum chamber. On the day of the experiment, the slides were fitted in a multiwell microarray hybridization cassette (ArrayIt, CA) to divide into eight wells. To each well, 400 μL of Ovalbumin (1% w/v, PBS) was added and rehydrated for 1 h in a humid chamber with gentle shaking. The solution was discarded followed by the addition of a 400 μL solution of the biotinylated PTOV B-subunit (30 μg mL<sup>-1</sup> dissolved in the same blocking buffer) in the individual well. The slides were then incubated for 2 h at ambient temperature with gentle shaking followed by washing with PBS-Tween (0.1% v/v) and PBS. The wells were then treated with Cy3-conjugated streptavidin as a secondary antibody (1:500 dilution in PBS). The slides were then incubated at ambient temperature for 1 h in a dark and humid chamber with gentle shaking. The slides were then washed, dried, and scanned using a Genepix 4000B scanner (Molecular Devices Corp., Union City, CA) at a wavelength of 532 nm. Data analysis was performed using the Genepix Pro 7.3 software (Molecular Devices Corp., Union City, CA).

**DMB-HPLC Analysis.** Sias were derivatized in a DMB reagent buffer containing a final concentration of 7 mM DMB, 9 mM sodium hydrosulfite, and 0.5 M of 2-mercaptoethanol added with either trifluoroacetic acid (20 mM) or acetic acid (0.8 M). The derivatized samples were analyzed on a LaChrom Elite HPLC (Hitachi) system with a Phenomenex Gemini C18 column (5 μm, 250 mm × 4.6 mm) at RT. A gradient elution (water/acetonitrile/methanol = 86:7:7 → 82:11:7) was used. The eluted analytes were detected with a fluorescence detector at an excitation of 373 nm and emission of 448 nm. The data collection time was extended to 75 min. For migration studies, the OAcSia monosaccharides or glycans were dissolved in a sodium acetate solution (50 mM) at various pHs (3.0–8.0). The starting material was diluted in the same volume of water as a control. All the samples were incubated in a 37 °C water bath for 2 h. The Sias were released from glycans by adding 10 mU of AUS and incubated at 37 °C for 10 min. The samples were then cooled on ice and derivatized with the DMB reagent containing acetic acid followed by HPLC analysis. The designation of Sia species was based on retention times compared to the Sia standards from bovine submaxillary mucin. The experiments were repeated three times showing similar results. For periodate treatment study, 50 μM sialoglycans was prepared in PBS (pH adjusted to 6.5) from a 10 mM stock. The sialoglycans were treated with 6 mM sodium periodate (2 M stock) at 20 °C for 2 h in the dark. Aliquots were taken every 30 min, and the generated aldehyde was reduced by the addition of sodium borohydride (1 M stock) to a final concentration of 30 mM at 20 °C for 20 min. The Sias and C7 analogues were released with 80 °C for 60 min. The sample was derivatized with DMB reagent containing acetic acid

followed by HPLC analysis. The relative peak area ratio of the C7-Neu5Ac analog to the nonperiodate treated Neu5Ac was applied to all samples for the calculation of the molar percentage of each Sia species.

**NMR Analysis of O-Acetylated Sia Monosaccharides.** Two samples of different concentrations and pHs were prepared. The first sample was prepared by dissolving 15 mg of O-acetylated Sia in 500  $\mu\text{L}$  of  $\text{D}_2\text{O}$  (80 mM). Its  $\text{pH}^*$  was adjusted to 3.5 (monitored by a micro pH meter) using HCl in  $\text{D}_2\text{O}$  solutions (100 mM, 10 mM, and 1 mM stock solutions prepared from 1 M HCl in  $\text{H}_2\text{O}$  by dilution with  $\text{D}_2\text{O}$ ).<sup>44,45</sup> It was then subjected to one-dimensional  $^1\text{H}$  NMR experiments performed on an 800 MHz Bruker Avance III spectrometer at the University of California, Davis.  $^1\text{H}$  NMR spectra were recorded at the beginning ( $t = 0$  min), every 10 min over the first 2 h, and at the end of the incubation period (12 h) at 310.15 K and  $\text{pH}^* = 3.5$ . The sample  $\text{pH}^*$  was checked again using a micro-pH meter at the end of the incubation and was shown to be stable. The  $\text{pH}^*$  of this sample was adjusted to 7.0 using NaOH in  $\text{D}_2\text{O}$  solutions (100 mM, 10 mM, and 1 mM stock solutions prepared by dissolving NaOH in  $\text{D}_2\text{O}$  followed by serial dilution with  $\text{D}_2\text{O}$ ), and the sample NMR experiments were carried out with  $\text{pH}^*$  rechecked at the end of the incubation. NMR data were processed and analyzed with MestreNova 12.0.4 software. Assignments of OAcSia in the  $^1\text{H}$  NMR study were verified by  $^1\text{H}$ - $^{13}\text{C}$  HSQC,<sup>22</sup> HSQC-TOCSY,<sup>24</sup> and LR-HSQMBC<sup>25</sup> experiments. For these experiments, a second sample was prepared by dissolving 20 mg of O-acetylated Sia in 450  $\mu\text{L}$  of  $\text{D}_2\text{O}$  (127 mM) at  $\text{pH}^* = 6.14$ . The 2D NMR experiments were carried out on a Bruker 700 MHz NMR AV III HD instrument equipped with a triple resonance, xyz gradient TCI cryoprobe at the FDA. Two-dimensional NMR spectra were acquired using Bruker's Topspin 3.6.1 software (<http://www.bruker.com>). The carrier frequency, spectral window, and number of points for the direct and indirect dimensions were set to 4.7 ppm, 60 ppm; 10 ppm, 86 ppm; and 2048, 1024, respectively, for the  $^1\text{H}$ - $^{13}\text{C}$  HSQC, HSQC-TOCSY, and LR-HSQMBC experiments. A DIPSI-2 mixing sequence with mixing times of 30 and 60 ms were used for the  $^1\text{H}$ - $^{13}\text{C}$  HSQC-TOCSY. The INEPT delay in the LR-HSQMBC was optimized for a 6 Hz long-range  $^1\text{H}$ - $^{13}\text{C}$  coupling constant. A multiplicity edited  $^1\text{H}$ - $^{13}\text{C}$  HSQC<sup>46</sup> was taken with 1024 points in the direct dimension and 2048 points in the indirect dimension using a nonuniform sampling data collection strategy in the indirect dimension with 50% of the points being collected based on the schedules from the Wagner group.<sup>47</sup> 2D NMR data were processed with NMRpipe,<sup>48</sup> using a square cosine bell to process both the direct and indirect dimensions. The direct dimension was referenced based on the DSS  $^1\text{H}$  signal, and  $^{13}\text{C}$  was referenced indirectly from the absolute DSS  $^1\text{H}$  frequency. Nonuniformly sampled multiplicity edited  $^1\text{H}$ - $^{13}\text{C}$  HSQC data were reconstructed using the SMILE reconstruction algorithm.<sup>49</sup>

**NMR Analysis of O-Acetylated Sialoglycans.**  $^1\text{H}$  NMR experiments of  $\text{pH}^*$ -dependent O-acetyl migration of O-acetylated sialoglycans were performed on an 800 MHz Bruker Avance III spectrometer at the University of California, Davis. The sample was prepared by dissolving 10 mg of Neu5,9Ac<sub>2</sub>Gal $\beta$ 4Glc $\beta$ ProN<sub>3</sub> in 500  $\mu\text{L}$  of  $\text{D}_2\text{O}$  (26 mM). Sample  $\text{pH}^*$  was adjusted to 3.5 by adding HCl in  $\text{D}_2\text{O}$ . It was incubated at 310.15 K for 2 h followed by adjusting the  $\text{pH}^*$  to 8.0 by adding NaOH in  $\text{D}_2\text{O}$  and incubation at 310.15 K for 2 h. The  $\text{pH}^*$  of the same sample was then adjusted back to 3.5. It was incubated at 310.15 K for 3 days followed by adjusting the  $\text{pH}^*$  to 8.0 and incubated at 310.15 K for 12 h.  $^1\text{H}$  NMR spectra were recorded at 0 min and at the end of each incubation period. Water suppression was applied when the spectra were recorded.  $^1\text{H}$  NMR data were processed and analyzed with TopSpin 3.6.1 software.

**LC-MS Analysis.** All mass spectrometry analyses were performed using an LTQ-XL Orbitrap Discovery (Thermo Scientific) mass spectrometer in the negative mode. LC profiling of sialic acid was done on a C18 column using an ion-pairing reagent in running buffer. The running buffer was constituted of dibutyl amine (DBA; 5 mM), acetic acid (8 mM), and methanol (70%). For the migration assay, deuterated OAc-Sia (almost all Neu5,9Ac<sub>2</sub>) was mixed with nondeuterated OAc-Sia to obtain a starting mixture containing

desired percentage of each OAcSia species (around 1 mM). The starting mixture (1.5  $\mu\text{L}$ ) was added to 18.5  $\mu\text{L}$  of 50 mM sodium acetate with a pH value in the range of 5.0–8.0 and incubated at 37  $^\circ\text{C}$ . Then, 4  $\mu\text{L}$  of the mixture was injected for LC-MS analysis. For the periodate treatment study, 50 mM of sialoglycans was prepared in PBS (pH adjusted to 6.5) from a 10 mM stock. The sialoglycans were treated with 6 mM sodium periodate (2 M stock) at 20  $^\circ\text{C}$  for 2 h in the dark. The resulting aldehyde was reduced by the addition of  $\text{NaBH}_4$  (1 M stock) to a final concentration of 30 mM at 20  $^\circ\text{C}$  for 20 min. The pH was adjusted to around 7.0 with 2 M acetic acid before injection for LC-MS analysis.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acscchembio.0c00998>.

NMR spectra and additional tables and figures (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Authors

**Ajit Varki** – Glycobiology Research and Training Center, Departments of Medicine and Cellular and Molecular Medicine, University of California, San Diego, California 92093, United States; Phone: +1 530-754-6037; Email: [xiichen@ucdavis.edu](mailto:xiichen@ucdavis.edu); Fax: +1 530-752-8995

**Xi Chen** – Department of Chemistry, University of California, Davis, California 95616, United States; [orcid.org/0000-0002-3160-614X](https://orcid.org/0000-0002-3160-614X); Phone: +1 858-534-2214; Email: [alvarki@ucsd.edu](mailto:alvarki@ucsd.edu); Fax: +1 858-534-5611

### Authors

**Yang Ji** – Glycobiology Research and Training Center, Departments of Medicine and Cellular and Molecular Medicine, University of California, San Diego, California 92093, United States

**Aniruddha Sasmal** – Glycobiology Research and Training Center, Departments of Medicine and Cellular and Molecular Medicine, University of California, San Diego, California 92093, United States

**Wanqing Li** – Department of Chemistry, University of California, Davis, California 95616, United States

**Lisa Oh** – Department of Chemistry, University of California, Davis, California 95616, United States; [orcid.org/0000-0001-8572-9738](https://orcid.org/0000-0001-8572-9738)

**Saurabh Srivastava** – Glycobiology Research and Training Center, Departments of Medicine and Cellular and Molecular Medicine, University of California, San Diego, California 92093, United States

**Audra A. Hargett** – Laboratory of Bacterial Polysaccharides, Food and Drug Administration (FDA), Silver Spring, Maryland 20993, United States

**Brian R. Wasik** – Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, New York 14853, United States; [orcid.org/0000-0001-5442-3883](https://orcid.org/0000-0001-5442-3883)

**Hai Yu** – Department of Chemistry, University of California, Davis, California 95616, United States; [orcid.org/0000-0002-4378-0532](https://orcid.org/0000-0002-4378-0532)

**Sandra Diaz** – Glycobiology Research and Training Center, Departments of Medicine and Cellular and Molecular Medicine, University of California, San Diego, California 92093, United States

**Biswa Choudhury** – Glycobiology Research and Training Center, Departments of Medicine and Cellular and Molecular



Medicine, University of California, San Diego, California 92093, United States

Colin R. Parrish – Baker Institute for Animal Health, College of Veterinary Medicine, Cornell University, Ithaca, New York 14853, United States; [orcid.org/0000-0002-1836-6655](https://orcid.org/0000-0002-1836-6655)

Darón I. Freedberg – Laboratory of Bacterial Polysaccharides, Food and Drug Administration (FDA), Silver Spring, Maryland 20993, United States; [orcid.org/0000-0003-1991-5174](https://orcid.org/0000-0003-1991-5174)

Lee-Ping Wang – Department of Chemistry, University of California, Davis, California 95616, United States; [orcid.org/0000-0003-3072-9946](https://orcid.org/0000-0003-3072-9946)

Complete contact information is available at:  
<https://pubs.acs.org/10.1021/acscchembio.0c00998>

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

This work was supported by the United States National Institutes of Health grant no. R01AI130684 (to X.C. and A.V.) and grant no. R01GM32373 (to A.V.). We thank A. Lewis for revising and providing comments on the manuscript.

## ABBREVIATIONS

Ac, acetyl group; AUS, *Arthrobacter ureafaciens* sialidase; BSM, bovine submaxillary mucin; CPS, capsular polysaccharide; DMB, 1,2-diamino-4,5-methylenedioxybenzene; ESI, electrospray ionization; HE, hemagglutinin-esterase; HPLC, high performance liquid chromatography; LC-ESI-MS, liquid chromatography-electrospray ionization-mass spectrometry; MS, mass spectrometry; Neu5Ac, *N*-acetylneuraminic acid; Neu5,9Ac<sub>2</sub>, 9-*O*-acetyl-*N*-acetylneuraminic acid; Neu5,8Ac<sub>2</sub>, 8-*O*-acetyl-*N*-acetylneuraminic acid; Neu5,7Ac<sub>2</sub>, 7-*O*-acetyl-*N*-acetylneuraminic acid; Neu5Ac9NAc, 9-acetamido-9-deoxy-*N*-acetylneuraminic acid; NMR, nuclear magnetic resonance; OAc, *O*-acetyl group; PToV, esterase-inactivated porcine torovirus; Sia, sialic acid; XIC, extracted ion current

## REFERENCES

- (1) Varki, A., Schnaar, R. L., and Schauer, R. (2015) *Essentials of Glycobiology*, pp 179–195, Cold Spring Harbor Laboratory Press.
- (2) Schauer, R., and Kamerling, J. P. (2018) Exploration of the sialic acid world. *Adv. Carbohydr. Chem. Biochem.* 75, 1–213.
- (3) Deng, L., Chen, X., and Varki, A. (2013) Exploration of sialic acid diversity and biology using sialoglycan microarrays. *Biopolymers* 99, 650–665.
- (4) Varki, N. M., and Varki, A. (2007) Diversity in cell surface sialic acid presentations: implications for biology and disease. *Lab. Invest.* 87, 851–857.
- (5) Muñoz-Barroso, I., García-Sastre, A., Villar, E., Manuguerra, J. C., Hannoun, C., and Cabezas, J. A. (1992) Increased influenza A virus sialidase activity with *N*-acetyl-9-*O*-acetylneuraminic acid-containing substrates resulting from influenza C virus *O*-acetyltransferase action. *Virus Res.* 25, 145–153.
- (6) Muchmore, E. A., and Varki, A. (1987) Selective inactivation of influenza C esterase: a probe for detecting 9-*O*-acetylated sialic acids. *Science* 236, 1293–1295.
- (7) Song, H., Qi, J., Khedri, Z., Diaz, S., Yu, H., Chen, X., Varki, A., Shi, Y., and Gao, G. F. (2016) An open receptor-binding cavity of hemagglutinin-esterase-fusion glycoprotein from newly-identified influenza D virus: Basis for its broad cell tropism. *PLoS Pathog.* 12, No. e1005411–e1005411.
- (8) Barnard, K. N., Wasik, B. R., LaClair, J. R., Buchholz, D. W., Weichert, W. S., Alford-Lawrence, B. K., Aguilar, H. C., and Parrish, C. R. (2019) Expression of 9-*O*- and 7,9-*O*-acetyl modified sialic acid in cells and their effects on influenza viruses. *mBio* 10, e02490-19.
- (9) Liu, R., Sreenivasan, C., Yu, H., Sheng, Z., Newkirk, S. J., An, W., Smith, D. F., Chen, X., Wang, D., and Li, F. (2020) Influenza D virus diverges from its related influenza C virus in the recognition of 9-*O*-acetylated *N*-acetyl- or *N*-glycolyl-neuraminic acid-containing glycan receptors. *Virology* 545, 16–23.
- (10) Weiman, S., Dahesh, S., Carlin, A. F., Varki, A., Nizet, V., and Lewis, A. L. (2009) Genetic and biochemical modulation of sialic acid *O*-acetylation on group B Streptococcus: phenotypic and functional impact. *Glycobiology* 19, 1204–1213.
- (11) Park, S. S. (2019) Post-glycosylation modification of sialic acid and its role in virus pathogenesis. *Vaccines* 7, 171.
- (12) Varki, A., and Diaz, S. (1984) The release and purification of sialic acids from glycoconjugates: Methods to minimize the loss and migration of *O*-acetyl groups. *Anal. Biochem.* 137, 236–247.
- (13) Kamerling, J. P., Schauer, R., Shukla, A. K., Stoll, S., Van Halbeek, H., and Vliegthart, J. F. (1987) Migration of *O*-acetyl groups in *N,O*-acetylneuraminic acids. *Eur. J. Biochem.* 162, 601–607.
- (14) Lemerrier, X., and Jones, C. (1996) Full <sup>1</sup>H NMR assignment and detailed *O*-acetylation patterns of capsular polysaccharides from *Neisseria meningitidis* used in vaccine production. *Carbohydr. Res.* 296, 83–96.
- (15) Lewis, A. L., Nizet, V., and Varki, A. (2004) Discovery and characterization of sialic acid *O*-acetylation in group B Streptococcus. *Proc. Natl. Acad. Sci. U. S. A.* 101, 11123–11128.
- (16) Diaz, S., Higa, H. H., and Varki, A. (1989) Glycoprotein sialate 7(9)-*O*-acetyltransferase from rat liver Golgi vesicles. *Methods Enzymol.* 179, 416–422.
- (17) Robinson, L. S., Lewis, W. G., and Lewis, A. L. (2017) The sialate *O*-acetyltransferase EstA from gut Bacteroidetes species enables sialidase-mediated cross-species foraging of 9-*O*-acetylated sialoglycans. *J. Biol. Chem.* 292, 11861–11872.
- (18) Lewis, A. L., Cao, H., Patel, S. K., Diaz, S., Ryan, W., Carlin, A. F., Thon, V., Lewis, W. G., Varki, A., Chen, X., and Nizet, V. (2007) NeuA sialic acid *O*-acetyltransferase activity modulates *O*-acetylation of capsular polysaccharide in group B Streptococcus. *J. Biol. Chem.* 282, 27562–27571.
- (19) Samraj, A. N., Pearce, O. M. T., Läubli, H., Crittenden, A. N., Bergfeld, A. K., Banda, K., Gregg, C. J., Bingman, A. E., Secret, P., Diaz, S. L., Varki, N. M., and Varki, A. (2015) A red meat-derived glycan promotes inflammation and cancer progression. *Proc. Natl. Acad. Sci. U. S. A.* 112, 542–547.
- (20) Klein, A., Diaz, S., Ferreira, I., Lamblin, G., Roussel, P., and Manzi, A. E. (1997) New sialic acids from biological sources identified by a comprehensive and sensitive approach: liquid chromatography-electrospray ionization-mass spectrometry (LC-ESI-MS) of SIA quinoxalinones. *Glycobiology* 7, 421–432.
- (21) Yu, H., Cheng, J., Ding, L., Khedri, Z., Chen, Y., Chin, S., Lau, K., Tiwari, V. K., and Chen, X. (2009) Chemoenzymatic synthesis of GD3 oligosaccharides and other disialyl glycans containing natural and non-natural sialic acids. *J. Am. Chem. Soc.* 131, 18467–18477.
- (22) Bodenhausen, G., and Ruben, D. J. (1980) Natural abundance nitrogen-15 NMR by enhanced heteronuclear spectroscopy. *Chem. Phys. Lett.* 69, 185–189.
- (23) Lerner, L., and Bax, A. (1986) Sensitivity-enhanced two-dimensional heteronuclear relayed coherence transfer NMR spectroscopy. *J. Magn. Reson.* (1969-1992) 69, 375–380.
- (24) Lerner, L., and Bax, A. (1987) Application of new, high-sensitivity, <sup>1</sup>H-<sup>13</sup>C-N.M.R.-spectral techniques to the study of oligosaccharides. *Carbohydr. Res.* 166, 35–46.
- (25) Williamson, R. T., Buevich, A. V., Martin, G. E., and Parella, T. (2014) LR-HSQMBC: A sensitive NMR technique to probe very long-range heteronuclear coupling pathways. *J. Org. Chem.* 79, 3887–3894.

- (26) Lo, L.-C., Hsiao, K.-F., Ueng, S.-H., and Wu, S.-H. (1999) Lipase-catalyzed acetylation of *N*-acetylneuraminic acid derivative. *Bioorg. Med. Chem. Lett.* 9, 709–712.
- (27) Haverkamp, J., van Halbeek, H., Dorland, L., Vliegthart, J. F., Pfeil, R., and Schauer, R. (1982) High-resolution <sup>1</sup>H-NMR spectroscopy of free and glycosidically linked *O*-acetylated sialic acids. *Eur. J. Biochem.* 122, 305–311.
- (28) Cheng, J., Yu, H., Lau, K., Huang, S., Chokhawala, H. A., Li, Y., Tiwari, V. K., and Chen, X. (2008) Multifunctionality of *Campylobacter jejuni* sialyltransferase CstII: Characterization of GD3/GT3 oligosaccharide synthase, GD3 oligosaccharide sialidase, and trans-sialidase activities. *Glycobiology* 18, 686–697.
- (29) Li, W., Battistel, M. D., Reeves, H., Oh, L., Yu, H., Chen, X., Wang, L. P., and Freedberg, D. I. (2020) A combined NMR, MD and DFT conformational analysis of 9-*O*-acetyl sialic acid-containing GM3 ganglioside glycan and its 9-*N*-acetyl mimic. *Glycobiology* 30, 787–801.
- (30) Khedri, Z., Xiao, A., Yu, H., Landig, C. S., Li, W., Diaz, S., Wasik, B. R., Parrish, C. R., Wang, L. P., Varki, A., and Chen, X. (2017) A chemical biology solution to problems with studying biologically important but unstable 9-*O*-acetyl sialic acids. *ACS Chem. Biol.* 12, 214–224.
- (31) Li, W., Xiao, A., Li, Y., Yu, H., and Chen, X. (2017) Chemoenzymatic synthesis of Neu5Ac9NAc-containing  $\alpha$ 2–3- and  $\alpha$ 2–6-linked sialosides and their use for sialidase substrate specificity studies. *Carbohydr. Res.* 451, 51–58.
- (32) Langereis, M. A., Bakkers, M. J.G., Deng, L., Padler-Karavani, V., Vervoort, S. J., Hulswit, R. J.G., van Vliet, A. L.W., Gerwig, G. J., de Poot, S. A.H., Boot, W., van Ederen, A. M., Heesters, B. A., van der Loos, C. M., van Kuppeveld, F. J.M., Yu, H., Huizinga, E. G., Chen, X., Varki, A., Kamerling, J. P., and de Groot, R. J. (2015) Complexity and diversity of the mammalian sialome revealed by nidovirus virolectins. *Cell Rep.* 11, 1966–1978.
- (33) Bobbitt, J. M. (1956) Periodate oxidation of carbohydrates, in *Advances in Carbohydrate Chemistry*, vol 11, pp 1–41, Elsevier.
- (34) Van Lenten, L., and Ashwell, G. (1971) Studies on the chemical and enzymatic modification of glycoproteins. A general method for the tritiation of sialic acid-containing glycoproteins. *J. Biol. Chem.* 246, 1889–1894.
- (35) Gahmberg, C. G., and Andersson, L. C. (1977) Selective radioactive labeling of cell surface sialoglycoproteins by periodate-tritiated borohydride. *J. Biol. Chem.* 252, 5888–5894.
- (36) Ramya, T. N. C., Weerapana, E., Cravatt, B. F., and Paulson, J. C. (2013) Glycoproteomics enabled by tagging sialic acid- or galactose-terminated glycans. *Glycobiology* 23, 211–221.
- (37) McCombs, J. E., and Kohler, J. J. (2016) Pneumococcal neuraminidase substrates identified through comparative proteomics enabled by chemoselective labeling. *Bioconjugate Chem.* 27, 1013–1022.
- (38) Tasnima, N., Yu, H., Yan, X., Li, W., Xiao, A., and Chen, X. (2019) Facile chemoenzymatic synthesis of Lewis a (Le(a)) antigen in gram-scale and sialyl Lewis a (sLe(a)) antigens containing diverse sialic acid forms. *Carbohydr. Res.* 472, 115–121.
- (39) Sato, C., Inoue, S., Matsuda, T., and Kitajima, K. (1998) Development of a highly sensitive chemical method for detecting  $\alpha$ 2→8-linked oligo/polysialic acid residues in glycoproteins blotted on the membrane. *Anal. Biochem.* 261, 191–197.
- (40) Sklarz, B. (1967) Organic chemistry of periodates. *Q. Rev., Chem. Soc.* 21, 3–28.
- (41) Wasik, B. R., Barnard, K. N., Ossiboff, R. J., Khedri, Z., Feng, K. H., Yu, H., Chen, X., Perez, D. R., Varki, A., and Parrish, C. R. (2017) Distribution of *O*-acetylated sialic acids among target host tissues for influenza virus. *mSphere* 2, No. e00379–00316.
- (42) Meng, C., Sasmal, A., Zhang, Y., Gao, T., Liu, C.-C., Khan, N., Varki, A., Wang, F., and Cao, H. (2018) Chemoenzymatic assembly of mammalian *O*-mannose glycans. *Angew. Chem., Int. Ed.* 57, 9003–9007.
- (43) Lu, N., Ye, J., Cheng, J., Sasmal, A., Liu, C.-C., Yao, W., Yan, J., Khan, N., Yi, W., Varki, A., and Cao, H. (2019) Redox-controlled site-specific  $\alpha$ 2–6-sialylation. *J. Am. Chem. Soc.* 141, 4547–4552.
- (44) Covington, A. K., Paabo, M., Robinson, R. A., and Bates, R. G. (1968) Use of the glass electrode in deuterium oxide and the relation between the standardized pD (paD) scale and the operational pH in heavy water. *Anal. Chem.* 40, 700–706.
- (45) Krezel, A., and Bal, W. (2004) A formula for correlating pKa values determined in D<sub>2</sub>O and H<sub>2</sub>O. *J. Inorg. Biochem.* 98, 161–166.
- (46) Sakhaei, P., and Bermel, W. (2015) A different approach to multiplicity-edited heteronuclear single quantum correlation spectroscopy. *J. Magn. Reson.* 259, 82–86.
- (47) Hyberts, S. G., Robson, S. A., and Wagner, G. (2013) Exploring signal-to-noise ratio and sensitivity in non-uniformly sampled multidimensional NMR spectra. *J. Biomol. NMR* 55, 167–178.
- (48) Delaglio, F., Grzesiek, S., Vuister, G. W., Zhu, G., Pfeifer, J., and Bax, A. (1995) NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J. Biomol. NMR* 6, 277–293.
- (49) Ying, J., Delaglio, F., Torchia, D. A., and Bax, A. (2017) Sparse multidimensional iterative lineshape-enhanced (SMILE) reconstruction of both non-uniformly sampled and conventional NMR data. *J. Biomol. NMR* 68, 101–118.